

CONTRACT NO. HY/2012/07

**Hong Kong-Zhuhai-Macao Bridge Tuen Mun – Chek Lap Kok Link
(Southern Connection Viaduct Section)
Dolphin Quarterly Monitoring**

*10th Quarterly Progress Report (March-May 2016)
submitted to Gammon Construction Limited*

Submitted by
Samuel K.Y. Hung, Ph.D., Hong Kong Cetacean Research Project

7 July 2016

1. Introduction

- 1.1. The Tuen Mun-Chek Lap Kok Link (TM-CLKL) comprises a 1.6 km long dual 2-lane viaduct section between the Hong Kong Boundary Crossing Facilities (HKBCF) and the North Lantau Highway and associated roads at Tai Ho. Gammon Construction Limited (hereinafter called the “Contractor”) was awarded as the main contractor of “Contract No. HY/2012/07 – Hong Kong-Zhuhai-Macao Bridge Tuen Mun-Chek Lap Kok Link – Southern Connection Viaduct Section”.
- 1.2. According to the updated Environmental Monitoring and Audit (EM&A) Manual (for TM-CLKL), monthly line-transect vessel surveys for Chinese White Dolphin should be conducted to cover the Northwest (NWL) and Northeast Lantau (NEL) survey areas as in AFCD annual marine mammal monitoring programme. However, as such surveys have been undertaken by the HKLR03 and HKBCF projects in the same areas (i.e. NWL and NEL), a combined monitoring approach is recommended by the Highways Department, that the TM-CLKL EM&A project can utilize the monitoring data collected by HKLR03 or HKBCF project to avoid any redundancy in monitoring effort. Such exemption for the dolphin monitoring will end upon the completion of the dolphin monitoring carried out by HKLR03 contract as well as the TM-CLKL Northern Connection Sub-Sea Tunnel Section (HY/2012/08)
- 1.3. In November 2013, the Director of Hong Kong Cetacean Research Project (HKCRP), Dr. Samuel Hung, has been appointed by Gammon Construction Limited as the dolphin specialist for the TM-CLKL Southern Viaduct Section EM&A project. He is responsible for the dolphin monitoring study, including the data collection on Chinese White Dolphins during the construction phase (i.e. impact period) of the TM-CLKL project in Northwest Lantau (NWL) and Northeast Lantau (NEL) survey areas.
- 1.4. During the construction period of HKLR, the dolphin specialist would be in charge of

reviewing and collating information collected by the HKLR03 dolphin monitoring programme to examine any potential impacts of TM-CLKL construction works on the dolphins.

- 1.5. From the monitoring results, any changes in dolphin occurrence within the study area will be examined for possible causes, and appropriate actions and additional mitigation measures will be recommended as necessary.
- 1.6. This report is the tenth quarterly progress report under the TM-CLKL construction phase dolphin monitoring programme submitted to the Gammon Construction Limited, summarizing the results of the surveys findings during the period of March to May 2016, utilizing the survey data collected by HKLR03 project.

2. Monitoring Methodology

2.1. Vessel-based Line-transect Survey

- 2.1.1. According to the requirement of the updated EM&A manual, dolphin monitoring programme should cover all transect lines in NEL and NWL survey areas (see Figure 1) twice per month throughout the entire construction period. The co-ordinates of all transect lines are shown in Table 1. The coordinates of several starting points have been revised due to the obstruction of the permanent structures in association to the construction works of HKLR and the southern viaduct of TM-CLKL, as well as provision of adequate buffer distance from the Airport Restricted Areas. The EPD issued a memo and confirmed that they had no objection on the revised transect lines on 19 August 2015, and the revised coordinates are in red and marked with an asterisk in Table 1.

Table 1 Co-ordinates of transect lines conducted by HKLR03 project

| Line No. | | Easting | Northing | | Line No. | Easting | Northing |
|----------|-------------|---------|----------|--|----------|---------|----------|
| 1 | Start Point | 804671 | 815456* | | 13 | 816506 | 819480 |
| 1 | End Point | 804671 | 831404 | | 13 | 816506 | 824859 |
| 2 | Start Point | 805475 | 815913* | | 14 | 817537 | 820220 |
| 2 | End Point | 805477 | 826654 | | 14 | 817537 | 824613 |
| 3 | Start Point | 806464 | 819435 | | 15 | 818568 | 820735 |
| 3 | End Point | 806464 | 822911 | | 15 | 818568 | 824433 |
| 4 | Start Point | 807518 | 819771 | | 16 | 819532 | 821420 |
| 4 | End Point | 807518 | 829230 | | 16 | 819532 | 824209 |
| 5 | Start Point | 808504 | 820220 | | 17 | 820451 | 822125 |
| 5 | End Point | 808504 | 828602 | | 17 | 820451 | 823671 |
| 6 | Start Point | 809490 | 820466 | | 18 | 821504 | 822371 |
| 6 | End Point | 809490 | 825352 | | 18 | 821504 | 823761 |

| | | | | | | | | |
|----|-------------|--------|---------|--|----|-------------|--------|--------|
| 7 | Start Point | 810499 | 820880* | | 19 | Start Point | 822513 | 823268 |
| 7 | End Point | 810499 | 824613 | | 19 | End Point | 822513 | 824321 |
| 8 | Start Point | 811508 | 821123* | | 20 | Start Point | 823477 | 823402 |
| 8 | End Point | 811508 | 824254 | | 20 | End Point | 823477 | 824613 |
| 9 | Start Point | 812516 | 821303* | | 21 | Start Point | 805476 | 827081 |
| 9 | End Point | 812516 | 824254 | | 21 | End Point | 805476 | 830562 |
| 10 | Start Point | 813525 | 820872 | | 22 | Start Point | 806464 | 824033 |
| 10 | End Point | 813525 | 824657 | | 22 | End Point | 806464 | 829598 |
| 11 | Start Point | 814556 | 818853* | | 23 | Start Point | 814559 | 821739 |
| 11 | End Point | 814556 | 820992 | | 23 | End Point | 814559 | 824768 |
| 12 | Start Point | 815542 | 818807 | | | | | |
| 12 | End Point | 815542 | 824882 | | | | | |

Note: Co-ordinates in red and marked with asterisk are revised co-ordinates of transect line.

- 2.1.2. The HKLR03 survey team used standard line-transect methods (Buckland et al. 2001) to conduct the systematic vessel surveys, and followed the same technique of data collection that has been adopted over the last 16 years of marine mammal monitoring surveys in Hong Kong developed by HKCRP (see Hung 2013, 2014). For each monitoring vessel survey, a 15-m inboard vessel with an open upper deck (about 4.5 m above water surface) was used to make observations from the flying bridge area.
- 2.1.3. Two experienced observers (a data recorder and a primary observer) made up the on-effort survey team, and the survey vessel transited different transect lines at a constant speed of 13-15 km per hour. The data recorder searched with unaided eyes and filled out the datasheets, while the primary observer searched for dolphins and porpoises continuously through 7 x 50 *Fujinon* marine binoculars. Both observers searched the sea ahead of the vessel, between 270° and 90° (in relation to the bow, which is defined as 0°). One to two additional experienced observers were available on the boat to work in shift (i.e. rotate every 30 minutes) in order to minimize fatigue of the survey team members. All observers were experienced in small cetacean survey techniques and identifying local cetacean species.
- 2.1.4. During on-effort survey periods, the survey team recorded effort data including time, positions (latitude and longitude), weather conditions (Beaufort sea state and visibility), and distance traveled in each series (a continuous period of search effort) with the assistance of a handheld GPS (*Garmin eTrex Legend*).
- 2.1.5. Data including time, position and vessel speed were also automatically and continuously logged by handheld GPS throughout the entire survey for subsequent review.
- 2.1.6. When dolphins were sighted, the survey team would end the survey effort, and immediately record the initial sighting distance and angle of the dolphin group from the survey vessel, as well as the sighting time and position. Then the research vessel was

diverted from its course to approach the animals for species identification, group size estimation, assessment of group composition, and behavioural observations. The perpendicular distance (PSD) of the dolphin group to the transect line was later calculated from the initial sighting distance and angle.

- 2.1.7. Survey effort being conducted along the parallel transect lines that were perpendicular to the coastlines (as indicated in Figure 1) was labeled as “primary” survey effort, while the survey effort conducted along the connecting lines between parallel lines was labeled as “secondary” survey effort. According to HKCRP long-term dolphin monitoring data, encounter rates of Chinese white dolphins deduced from effort and sighting data collected along primary and secondary lines were similar in NEL and NWL survey areas. Therefore, both primary and secondary survey effort were presented as on-effort survey effort in this report.

2.2. Photo-identification Work

- 2.2.1. When a group of Chinese White Dolphins were sighted during the line-transect survey, the HKLR03 survey team would end effort and approach the group slowly from the side and behind to take photographs of them. Every attempt was made to photograph every dolphin in the group, and even photograph both sides of the dolphins, since the colouration and markings on both sides may not be symmetrical.
- 2.2.2. A professional digital camera (*Canon* EOS 7D or 60D model), equipped with long telephoto lenses (100-400 mm zoom), were available on board for researchers to take sharp, close-up photographs of dolphins as they surfaced. The images were shot at the highest available resolution and stored on Compact Flash memory cards for downloading onto a computer.
- 2.2.3. All digital images taken in the field were first examined, and those containing potentially identifiable individuals were sorted out. These photographs would then be examined in greater detail, and were carefully compared to the existing Chinese White Dolphin photo-identification catalogue maintained by HKCRP since 1995.
- 2.2.4. Chinese White Dolphins can be identified by their natural markings, such as nicks, cuts, scars and deformities on their dorsal fin and body, and their unique spotting patterns were also used as secondary identifying features (Jefferson 2000).
- 2.2.5. All photographs of each individual were then compiled and arranged in chronological order, with data including the date and location first identified (initial sighting), re-sightings, associated dolphins, distinctive features, and age classes entered into a computer database.

2.3. Data Analysis

- 2.3.1. Distribution Analysis – The line-transect survey data was integrated with the Geographic Information System (GIS) in order to visualize and interpret different spatial and temporal

patterns of dolphin distribution using sighting positions. Location data of dolphin groups were plotted on map layers of Hong Kong using a desktop GIS (ArcView[®] 3.1) to examine their distribution patterns in details. The dataset was also stratified into different subsets to examine distribution patterns of dolphin groups with different categories of group sizes, young calves and activities.

- 2.3.2. Encounter rate analysis – Encounter rates of Chinese white dolphins (number of on-effort sightings per 100 km of survey effort, and total number of dolphins sighted on-effort per 100 km of survey effort) were calculated in NEL and NWL survey areas in relation to the amount of survey effort conducted during each month of monitoring survey. Only data collect under Beaufort 3 or below condition would be used for the encounter rate analyses. Dolphin encounter rates were calculated in two ways for comparisons with the HZMB baseline monitoring results as well as to AFCD long-term marine mammal monitoring results.

Firstly, for the comparison with the HZMB baseline monitoring results, the encounter rates were calculated using primary survey effort alone. The average encounter rate of sightings (STG) and average encounter rate of dolphins (ANI) were deduced based on the encounter rates from six events during the present quarter (i.e. six sets of line-transect surveys in North Lantau), which was also compared with the one deduced from the six events during the baseline period (i.e. six sets of line-transect surveys in North Lantau).

Secondly, the encounter rates were calculated using both primary and secondary survey effort collected under Beaufort 3 or below condition as in AFCD long-term monitoring study. The encounter rate of sightings and dolphins were deduced by dividing the total number of on-effort sightings (STG) and total number of dolphins (ANI) by the amount of survey effort for the present quarterly period.

- 2.3.3. Quantitative grid analysis on habitat use – To conduct quantitative grid analysis of habitat use, positions of on-effort sightings of Chinese White Dolphins collected during the quarterly impact phase monitoring period were plotted onto 1-km² grids among NWL and NEL survey areas on GIS. Sighting densities (number of on-effort sightings per km²) and dolphin densities (total number of dolphins from on-effort sightings per km²) were then calculated for each 1 km by 1 km grid with the aid of GIS. Sighting density grids and dolphin density grids were then further normalized with the amount of survey effort conducted within each grid. The total amount of survey effort spent on each grid was calculated by examining the survey coverage on each line-transect survey to determine how many times the grid was surveyed during the study period. For example, when the survey boat traversed through a specific grid 50 times, 50 units of survey effort were counted for that grid. With the amount of survey effort calculated for each grid, the sighting density and dolphin density of each grid were then normalized (i.e. divided by the unit of survey effort).

The newly-derived unit for sighting density was termed SPSE, representing the number of on-effort sightings per 100 units of survey effort. In addition, the derived unit for actual dolphin density was termed DPSE, representing the number of dolphins per 100 units of survey effort. Among the 1-km² grids that were partially covered by land, the

percentage of sea area was calculated using GIS tools, and their SPSE and DPSE values were adjusted accordingly. The following formulae were used to estimate SPSE and DPSE in each 1-km² grid within the study area:

$$\text{SPSE} = ((S / E) \times 100) / \text{SA}\%$$

$$\text{DPSE} = ((D / E) \times 100) / \text{SA}\%$$

where S = total number of on-effort sightings
D = total number of dolphins from on-effort sightings
E = total number of units of survey effort
SA% = percentage of sea area

- 2.3.4. Behavioural analysis – When dolphins were sighted during vessel surveys, their behaviour was observed. Different activities were categorized (i.e. feeding, socializing, traveling, and milling/resting) and recorded on sighting datasheets. This data was then input into a separate database with sighting information, which can be used to determine the distribution of behavioural data with a desktop GIS. Distribution of sightings of dolphins engaged in different activities and behaviours would then be plotted on GIS and carefully examined to identify important areas for different activities of the dolphins.
- 2.3.5. Ranging pattern analysis – Location data of individual dolphins that occurred during the 3-month impact phase monitoring period were obtained from the dolphin sighting database and photo-identification catalogue. To deduce home ranges for individual dolphins using the fixed kernel methods, the program Animal Movement Analyst Extension, was loaded as an extension with ArcView[®] 3.1 along with another extension Spatial Analyst 2.0. Using the fixed kernel method, the program calculated kernel density estimates based on all sighting positions, and provided an active interface to display kernel density plots. The kernel estimator then calculated and displayed the overall ranging area at 95% UD level.

3. Monitoring Results

- 3.1. *Summary of survey effort and dolphin sightings*
- 3.1.1. During the period of March to May 2016, six sets of systematic line-transect vessel surveys were conducted under the HKLR03 monitoring works to cover all transect lines in NWL and NEL survey areas twice per month.
- 3.1.2. From these HKLR03 surveys, a total of 896.56 km of survey effort was collected, with 90.3% of the total survey effort being conducted under favourable weather conditions (i.e. Beaufort Sea State 3 or below with good visibility). Among the two areas, 341.16 km and 555.40 km of survey effort were conducted in NEL and NWL survey areas respectively.
- 3.1.3. The total survey effort conducted on primary lines was 657.94 km, while the effort on secondary lines was 238.62 km. Survey effort conducted on both primary and secondary lines were considered as on-effort survey data. A summary table of the survey effort is

shown in Appendix I.

- 3.1.4. During the six sets of HKLR03 monitoring surveys from March to May 2016, a total of seven groups of 22 Chinese White Dolphins were sighted. Four of the seven dolphin sightings were made during on-effort search, while three of the four on-effort dolphin sightings were made on primary lines. In this quarterly period, all dolphin groups were sighted in NWL, while none was sighted at all in NEL. A summary table of the dolphin sightings is shown in Appendix II.
- 3.2. *Distribution*
- 3.2.1. Distribution of dolphin sightings made during the HKLR03 monitoring surveys from March to May 2016 is shown in Figure 1. Dolphin sightings made in the present quarter were mostly located to the north of Lung Kwu Chau, while one sighting each was made to the southwest of Lung Kwu Chau and at the northeast corner of the airport platform respectively (Figure 1). Notably, four of the five sightings located to the north of Lung Kwu Chau were all made on the same survey day.
- 3.2.2. Notably, none of the dolphin sightings was made near the TM-CLKL alignment, HKLR09 alignment or HKLR03 reclamation site. On the other hand, one dolphin group was sighted near the HKBCF reclamation site (Figure 1).
- 3.2.3. Sighting distribution of dolphins during the present impact phase monitoring period (March to May 2016) was drastically different from the one during the baseline monitoring period (September to November 2011). In the present quarter, dolphins have disappeared from the NEL region, which was in stark contrast to their frequent occurrence around the Brothers Islands, near Shum Shui Kok and in the vicinity of HKBCF reclamation site during the baseline period (Figure 1). The nearly complete abandonment of NEL region by the dolphins has been consistently recorded in the past three years of HKLR03 monitoring, which has resulted in zero to extremely low dolphin encounter rates in this area.
- 3.2.4. In NWL survey area, dolphin occurrence was also very different between the baseline and the present impact phase periods. During the present impact monitoring period, much fewer dolphins occurred in this survey area (mostly near Lung Kwu Chau) than during the baseline period, when many dolphin groups were frequently sighted between Lung Kwu Chau and Black Point, around Sha Chau, near Pillar Point and to the west of the Chek Lap Kok Airport (Figure 1).
- 3.2.5. Another comparison in dolphin distribution was made between the four quarterly periods of spring months in 2013-16 (Figure 2). Among the four spring periods, dolphins were regularly sighted throughout the North Lantau region in 2013, but their usage there have been significantly reduced to a very low level in 2015 and 2016 (Figure 2).
- 3.3. *Encounter rate*
- 3.3.1. During the present quarterly period, the encounter rates of Chinese White Dolphins deduced from the survey effort and on-effort sighting data from the primary transect lines under favourable conditions (Beaufort 3 or below) for each set of the HKLR03 surveys in

HK CETACEAN RESEARCH PROJECT 香港鯨豚研究計劃

NEL and NWL are shown in Table 2. The average encounter rates deduced from the six sets of HKLR03 surveys were also compared with the ones deduced from the baseline monitoring period (September – November 2011) (Table 3).

Table 2. Dolphin encounter rates (sightings per 100 km of survey effort) during March to May 2016

| SURVEY AREA | DOLPHIN MONITORING DATES | Encounter rate (STG) (no. of on-effort dolphin sightings per 100 km of survey effort) | Encounter rate (ANI) (no. of dolphins from all on-effort sightings per 100 km of survey effort) |
|------------------|--------------------------|--|--|
| | | Primary Lines Only | Primary Lines Only |
| Northeast Lantau | Set 1 (7 & 11 Mar 2016) | 0.00 | 0.00 |
| | Set 2 (22 & 23 Mar 2016) | 0.00 | 0.00 |
| | Set 3 (5 & 12 Apr 2016) | 0.00 | 0.00 |
| | Set 4 (15 & 19 Apr 2016) | 0.00 | 0.00 |
| | Set 5 (3 & 12 May 2016) | 0.00 | 0.00 |
| | Set 6 (17 & 26 May 2016) | 0.00 | 0.00 |
| Northwest Lantau | Set 1 (7 & 11 Mar 2016) | 0.00 | 0.00 |
| | Set 2 (22 & 23 Mar 2016) | 1.59 | 4.78 |
| | Set 3 (5 & 12 Apr 2016) | 2.20 | 17.59 |
| | Set 4 (15 & 19 Apr 2016) | 2.10 | 6.31 |
| | Set 5 (3 & 12 May 2016) | 0.00 | 0.00 |
| | Set 6 (17 & 26 May 2016) | 0.00 | 0.00 |

Table 3. Comparison of average dolphin encounter rates from impact monitoring period (March to May 2016) and baseline monitoring period (September – November 2011) (Note: encounter rates deduced from the baseline monitoring period have been recalculated based only on survey effort and on-effort sighting data made along the primary transect lines under favourable conditions; \pm denotes the standard deviation of the average encounter rates)

| | Encounter rate (STG) (no. of on-effort dolphin sightings per 100 km of survey effort) | | Encounter rate (ANI) (no. of dolphins from all on-effort sightings per 100 km of survey effort) | |
|------------------|--|---------------------------|--|---------------------------|
| | March – May 2016 | September - November 2011 | March – May 2016 | September - November 2011 |
| Northeast Lantau | 0.0 | 6.00 \pm 5.05 | 0.0 | 22.19 \pm 26.81 |
| Northwest Lantau | 0.98 \pm 1.10 | 9.85 \pm 5.85 | 4.78 \pm 6.85 | 44.66 \pm 29.85 |

3.3.2. To facilitate the comparison with the AFCD long-term monitoring results, the encounter rates were also calculated for the present quarter using both primary and secondary survey effort. The encounter rates of sightings (STG) and dolphins (ANI) in NWL were 0.83 sightings and 3.54 dolphins per 100 km of survey effort respectively, while the encounter rates of sightings (STG) and dolphins (ANI) in NEL were both nil for this quarter.

3.3.3. In NEL, the average dolphin encounter rates (both STG and ANI) in the present three-month impact monitoring period were both zero with no sighting being made, and such extremely low occurrence of dolphins in NEL have been consistently recorded in the past thirteen quarters of HKLR03 monitoring (Table 4). This is a serious concern as the dolphin occurrence in NEL in the past few years (0.0-1.0 for ER(STG) and 0.0-3.9 for ER(ANI)) have been exceptionally low when compared to the baseline period (Table 4). Dolphins have been virtually absent from NEL waters since January 2014, with only two groups of five dolphins sighted there since then despite consistent and intensive survey effort being conducted in this survey area.

Table 4. Comparison of average dolphin encounter rates in Northeast Lantau survey area from all quarters of HKLR03 impact monitoring period and baseline monitoring period (September-November 2011) (Note: encounter rates deduced from the baseline monitoring period have been recalculated based only on survey effort and on-effort sighting data made along the primary transect lines under favourable conditions; the encounter rates in spring months were highlighted in blue; \pm denotes the standard deviation of the average encounter rates)

| | Encounter rate (STG) (no. of on-effort dolphin sightings per 100 km of survey effort) | Encounter rate (ANI) (no. of dolphins from all on-effort sightings per 100 km of survey effort) |
|---|---|---|
| September-November 2011 (Baseline) | 6.00 \pm 5.05 | 22.19 \pm 26.81 |
| December 2012-February 2013 (Impact) | 3.14 \pm 3.21 | 6.33 \pm 8.64 |
| March-May 2013 (Impact) | 0.42 \pm 1.03 | 0.42 \pm 1.03 |
| June-August 2013 (Impact) | 0.88 \pm 1.36 | 3.91 \pm 8.36 |
| September-November 2013 (Impact) | 1.01 \pm 1.59 | 3.77 \pm 6.49 |
| December 2013-February 2014 (Impact) | 0.45 \pm 1.10 | 1.34 \pm 3.29 |
| March-May 2014 (Impact) | 0.00 | 0.00 |
| June-August 2014 (Impact) | 0.42 \pm 1.04 | 1.69 \pm 4.15 |
| September-November 2014 (Impact) | 0.00 | 0.00 |
| December 2014-February 2015 (Impact) | 0.00 | 0.00 |
| March-May 2015 (Impact) | 0.00 | 0.00 |
| June-August 2015 (Impact) | 0.44 \pm 1.08 | 0.44 \pm 1.08 |
| September-November 2015 (Impact) | 0.00 | 0.00 |
| December 2015-February 2016 (Impact) | 0.00 | 0.00 |
| March-May 2016 (Impact) | 0.00 | 0.00 |

3.3.4. On the other hand, the average dolphin encounter rates (STG and ANI) in NWL during the present impact phase monitoring period (reductions of 90.1% and 89.3% respectively) were only small fractions of the ones recorded during the three-month baseline period, indicating a dramatic decline in dolphin usage of this survey area as well during the present impact phase period (Table 5).

Table 5. Comparison of average dolphin encounter rates in Northwest Lantau survey area from all quarters of HKLR03 impact monitoring period and baseline monitoring period (September-November 2011) (Note: encounter rates deduced from the baseline monitoring period have been recalculated based only on survey effort and on-effort sighting data made along the primary transect lines under favourable conditions; the encounter rates in spring months were highlighted in blue; \pm denotes the standard deviation of the average encounter rates)

| | Encounter rate (STG) (no. of on-effort dolphin sightings per 100 km of survey effort) | Encounter rate (ANI) (no. of dolphins from all on-effort sightings per 100 km of survey effort) |
|---|---|---|
| September-November 2011 (Baseline) | 9.85 \pm 5.85 | 44.66 \pm 29.85 |
| December 2012-February 2013 (Impact) | 8.36 \pm 5.03 | 35.90 \pm 23.10 |
| March-May 2013 (Impact) | 7.75 \pm 3.96 | 24.23 \pm 18.05 |
| June-August 2013 (Impact) | 6.56 \pm 3.68 | 27.00 \pm 18.71 |
| September-November 2013 (Impact) | 8.04 \pm 1.10 | 32.48 \pm 26.51 |
| December 2013-February 2014 (Impact) | 8.21 \pm 2.21 | 32.58 \pm 11.21 |
| March-May 2014 (Impact) | 6.51 \pm 3.34 | 19.14 \pm 7.19 |
| June-August 2014 (Impact) | 4.74 \pm 3.84 | 17.52 \pm 15.12 |
| September-November 2014 (Impact) | 5.10 \pm 4.40 | 20.52 \pm 15.10 |
| December 2014-February 2015 (Impact) | 2.91 \pm 2.69 | 11.27 \pm 15.19 |
| March-May 2015 (Impact) | 0.47 \pm 0.73 | 2.36 \pm 4.07 |
| June-August 2015 (Impact) | 2.53 \pm 3.20 | 9.21 \pm 11.57 |
| September-November 2015 (Impact) | 3.94 \pm 1.57 | 21.05 \pm 17.19 |
| December 2015-February 2016 (Impact) | 2.64 \pm 1.52 | 10.98 \pm 3.81 |
| March-May 2016 (Impact) | 0.98 \pm 1.10 | 4.78 \pm 6.85 |

- 3.3.5. During the same spring quarters, the dolphin encounter rates in NWL during the spring months of 2015 and 2016 were much lower than the ones recorded in spring months of 2013 and 2014 (Table 5). Such temporal trend should be closely monitored in the upcoming monitoring quarters.
- 3.3.6. As discussed recently in Hung (2015), the dramatic decline in dolphin usage of NEL waters in the past few years (including the declines in abundance, encounter rate and habitat use in NEL, as well as shifts of individual core areas and ranges away from NEL waters) was possibly related to the HZMB construction works that were commenced since 2012. It appeared that such noticeable decline has already extended to NWL waters progressively in the past few years.
- 3.3.7. A two-way ANOVA with repeated measures and unequal sample size was conducted to examine whether there were any significant differences in the average encounter rates

between the baseline and impact monitoring periods. The two variables that were examined included the two periods (baseline and impact phases) and two locations (NEL and NWL).

- 3.3.8. For the comparison between the baseline period and the present quarter (fourteenth quarter of the impact phase being assessed), the p-values for the differences in average dolphin encounter rates of STG and ANI were 0.0019 and 0.0173 respectively. If the alpha value is set at 0.05, significant differences were detected between the baseline and present quarters in both the average dolphin encounter rates of STG and ANI.
- 3.3.9. For the comparison between the baseline period and the cumulative quarters in impact phase (i.e. first fourteen quarters of the impact phase being assessed), the p-values for the differences in average dolphin encounter rates of STG and ANI were 0.000019 and 0.000005 respectively. Even if the alpha value is set at 0.00005, significant differences were still detected in both the average dolphin encounter rates of STG and ANI (i.e. between the two periods and the locations).
- 3.3.10. As indicated in both dolphin distribution patterns and encounter rates, dolphin usage has been significantly reduced in both NEL and NWL survey areas during the present quarterly period, and such low occurrence of dolphins has also been consistently documented in previous quarters. This raises serious concern, as the timing of the decline in dolphin usage in North Lantau waters coincided well with the construction schedule of the HZMB-related projects (Hung 2015).
- 3.3.11. To ensure the continuous usage of North Lantau waters by the dolphins, every possible measure should be implemented by the contractors and relevant authorities of HZMB-related works to minimize all disturbances to the dolphins.

3.4. Group size

- 3.4.1. Group size of Chinese White Dolphins ranged from one to eight individuals per group in North Lantau region during March to May 2016. The average dolphin group sizes from these three months were compared with the ones deduced from the baseline period in September to November 2011, as shown in Table 6.

Table 6. Comparison of average dolphin group sizes from impact monitoring period (March – May 2016) and baseline monitoring period (September – November 2011) (Note: \pm denotes the standard deviation of the average group size)

| | Average Dolphin Group Size | |
|-------------------------|----------------------------|---------------------------|
| | March – May 2016 | September – November 2011 |
| Overall | 3.14 \pm 2.27 (n = 7) | 3.72 \pm 3.13 (n = 66) |
| Northeast Lantau | N/A | 3.18 \pm 2.16 (n = 17) |
| Northwest Lantau | 3.14 \pm 2.27 (n = 7) | 3.92 \pm 3.40 (n = 49) |

- 3.4.2. The average dolphin group size in NWL waters during March to May 2016 was slightly

lower than the ones recorded during the three-month baseline period (Table 6). All except one dolphin groups were composed of 1-3 individuals only, while only one group was moderately large with eight individuals.

- 3.4.3. Distribution of the lone larger dolphin group (i.e. five individuals or more per group) during the present quarter is shown in Figure 3, with comparison to the one in baseline period. During the spring months of 2016, the only larger dolphin group was sighted to the southwest of Lung Kwu Chau (Figure 3). Such distribution pattern was very different from the baseline period, when the larger dolphin groups were more frequently sighted and more evenly distributed in NWL waters, with a few more sighted in NEL waters (Figure 3).
- 3.5. *Habitat use*
- 3.5.1. From March to May 2016, the only area being utilized by Chinese White Dolphins was to the north and southwest of Lung Kwu Chau (Figures 4a and 4b). All grids near TMCLKL/HKLR09 alignments as well as HKLR03/HKBCF reclamation sites did not record any presence of dolphins at all during on-effort search in the present quarterly period (Figure 4b).
- 3.5.2. It should be emphasized though that the amount of survey effort collected in each grid during the three-month period was fairly low (6-12 units of survey effort for most grids), and therefore the habitat use pattern derived from the three-month dataset should be treated with caution. A more complete picture of dolphin habitat use pattern should be examined when more survey effort for each grid will be collected throughout the impact phase monitoring programme.
- 3.5.3. When compared with the habitat use patterns during the baseline period, dolphin usage in NEL and NWL has drastically diminished in both areas during the present impact monitoring period (Figure 5). During the baseline period, many grids between Siu Mo To and Shum Shui Kok in NEL recorded moderately high to high dolphin densities, which was in stark contrast to the complete absence of dolphins there during the present impact phase period (Figure 5).
- 3.5.4. The density patterns were also very different in NWL between the baseline and impact phase monitoring periods, with higher dolphin usage throughout the area, especially around Sha Chau, near Black Point, to the west of the airport, as well as between Pillar Point and airport platform during the baseline period. In contrast, the only area with moderate to high dolphin densities was restricted to the waters near Lung Kwu Chau during the present impact phase period (Figure 5).
- 3.6. *Mother-calf pairs*
- 3.6.1. During the present quarterly period, neither unspotted calf nor unspotted juvenile was sighted with any female in the North Lantau region.
- 3.6.2. The absence of young calves in the present quarter was in stark contrast to their regular occurrence in North Lantau waters during the baseline period. This should be of a serious concern, and the occurrence of young calves in North Lantau waters should be

closely monitored in the upcoming quarters.

3.7. *Activities and associations with fishing boats*

- 3.7.1. Only one of the seven dolphin groups were engaged in feeding activity, while none of them was engaged in socializing, traveling or milling/resting activity during the three-month study period.
- 3.7.2. The percentage of sightings associated with feeding activities (14.3%) was similar to the one recorded during the baseline period (11.6%). However, it should be noted the sample sizes on total numbers of dolphin sightings during the present quarter (seven dolphin groups) was much lower than the baseline period (66 dolphin groups).
- 3.7.3. Distribution of dolphins engaged in various activities during the present impact phase period and the baseline period is shown in Figure 6. The only dolphin group engaged in feeding activity was sighted to the north of Lung Kwu Chau during the present quarterly period, which was very different from the baseline period when various dolphin activities occurred throughout the North Lantau region (Figure 6).
- 3.7.4. As consistently recorded in the past monitoring quarters, none of the seven dolphin groups was found to be associated with any operating fishing vessel in North Lantau waters during the present impact phase period.
- 3.8. *Summary of photo-identification works*
- 3.8.1. From March to May 2016, over 800 digital photographs of Chinese White Dolphins were taken during the impact phase monitoring surveys for the photo-identification work.
- 3.8.2. In total, 16 individuals sighted 18 times altogether were identified (see summary table in Appendix III and photographs of identified individuals in Appendix IV). All of these re-sightings were made in NWL.
- 3.8.3. The majority of identified individuals were sighted only once during the three-month period, with the exception of two individuals (NL48 and NL285) being sighted twice in the present quarter.
- 3.8.4. Notably, two of these 16 individuals (NL123 and NL320) were also sighted in West Lantau waters during the HKLR09 monitoring surveys from March to May 2016.
- 3.9. *Individual range use*
- 3.9.1. Ranging patterns of the 16 individuals identified during the three-month study period were determined by fixed kernel method, and are shown in Appendix V.
- 3.9.2. All identified dolphins sighted in the present quarter were utilizing NWL waters only, but have completely avoided NEL waters where many of them have utilized as their core areas in the past (Appendix V). This is in contrary to the extensive movements between NEL and NWL survey areas observed in the earlier impact monitoring quarters as well as the baseline period.

- 3.9.3. On the other hand, two individuals (NL123 and NL320) consistently utilized both North Lantau waters in the past have extended their range use to WL waters during the present quarter. In the upcoming quarters, individual range use and movements should be continuously monitored to examine whether there has been any consistent shifts of individual home ranges from North Lantau to West or Southwest Lantau, as such shift could possibly be related to the HZMB-related construction works (see Hung 2015).

4. Conclusion

- 4.1. During this quarter of dolphin monitoring, no adverse impact from the activities of the TMCLKL construction project on Chinese White Dolphins was noticeable from general observations.
- 4.2. Although the dolphins infrequently occurred along the alignment of TMCLKL southern connection viaduct in the past and during the baseline monitoring period, it is apparent that dolphin usage has been significantly reduced in NEL, and many individuals have shifted away from the important habitat around the Brothers Islands.
- 4.3. It is critical to monitor the dolphin usage in North Lantau region in the upcoming quarters, to determine whether the dolphins are continuously affected by the various construction activities in relation to the HZMB-related works, and whether suitable mitigation measure can be applied to revert the situation.

5. References

- Buckland, S. T., Anderson, D. R., Burnham, K. P., Laake, J. L., Borchers, D. L., and Thomas, L. 2001. Introduction to distance sampling: estimating abundance of biological populations. Oxford University Press, London.
- Hung, S. K. 2013. Monitoring of Marine Mammals in Hong Kong waters: final report (2012-13). An unpublished report submitted to the Agriculture, Fisheries and Conservation Department, 168 pp.
- Hung, S. K. 2014. Monitoring of marine mammals in Hong Kong waters – data collection: final report (2013-14). An unpublished report submitted to the Agriculture, Fisheries and Conservation Department of Hong Kong SAR Government, 231 pp.
- Hung, S. K. 2015. Monitoring of marine mammals in Hong Kong waters – data collection: final report (2014-15). An unpublished report submitted to the Agriculture, Fisheries and Conservation Department of Hong Kong SAR Government, 198 pp.
- Jefferson, T. A. 2000. Population biology of the Indo-Pacific hump-backed dolphin in Hong Kong waters. Wildlife Monographs 144:1-65.





Figure 1. Distribution of Chinese white dolphin sighting in Northwest and Northeast Lantau during HKLR03 impact phase (top) and baseline monitoring surveys (bottom)







Figure 2. Distribution of Chinese white dolphin sightings in Northwest and Northeast Lantau during the same spring quarters (March-May) of HKLR03 impact phase in 2013-16





Figure 3. Distribution of Chinese white dolphins with larger group sizes during HKLR03 impact phase (top) and baseline monitoring surveys (bottom) (green dots: group sizes of 5 or more; purple dots: group sizes of 10 or more)




Figure 4a. Sighting density of Chinese white dolphins with corrected survey effort per km² in Northeast and Northwest Lantau survey areas, using data collected during HKLR03 impact monitoring period (March-May 2016) (SPSE = no. of on-effort sightings per 100 units of survey effort)




Figure 4b. Density of Chinese white dolphins with corrected survey effort per km² in Northeast and Northwest Lantau survey areas, using data collected during HKLR03 impact monitoring period (March-May 2016) (DPSE = no. of dolphins per 100 units of survey effort)




Figure 5. Comparison of density of Chinese white dolphins with corrected survey effort per km² in Northwest and Northeast Lantau survey area between the impact monitoring period (March-May 2016) and baseline monitoring period (September-November 2011) (DPSE = no. of dolphins per 100 units of survey effort)





Figure 6. Distribution of Chinese white dolphins engaged in feeding (purple dots), socializing (pink dots) and traveling (green dots) activities during HKLR03 impact phase (top) and baseline monitoring surveys (bottom)

Appendix I. HKLR03 Survey Effort Database (March-May 2016)

(Abbreviations: BEAU = Beaufort Sea State; P = Primary Line Effort; S = Secondary Line Effort)

| DATE | AREA | BEAU | EFFORT | SEASON | VESSEL | TYPE | P/S |
|-----------|-----------|------|--------|--------|---------------|------|-----|
| 7-Mar-16 | NW LANTAU | 1 | 18.42 | SPRING | STANDARD31516 | HKLR | P |
| 7-Mar-16 | NW LANTAU | 2 | 10.78 | SPRING | STANDARD31516 | HKLR | P |
| 7-Mar-16 | NW LANTAU | 3 | 10.30 | SPRING | STANDARD31516 | HKLR | P |
| 7-Mar-16 | NW LANTAU | 1 | 2.50 | SPRING | STANDARD31516 | HKLR | S |
| 7-Mar-16 | NW LANTAU | 2 | 3.70 | SPRING | STANDARD31516 | HKLR | S |
| 7-Mar-16 | NW LANTAU | 3 | 6.70 | SPRING | STANDARD31516 | HKLR | S |
| 7-Mar-16 | NE LANTAU | 2 | 16.44 | SPRING | STANDARD31516 | HKLR | P |
| 7-Mar-16 | NE LANTAU | 2 | 10.46 | SPRING | STANDARD31516 | HKLR | S |
| 11-Mar-16 | NW LANTAU | 2 | 15.40 | SPRING | STANDARD31516 | HKLR | P |
| 11-Mar-16 | NW LANTAU | 3 | 16.20 | SPRING | STANDARD31516 | HKLR | P |
| 11-Mar-16 | NW LANTAU | 2 | 7.60 | SPRING | STANDARD31516 | HKLR | S |
| 11-Mar-16 | NW LANTAU | 3 | 0.30 | SPRING | STANDARD31516 | HKLR | S |
| 11-Mar-16 | NE LANTAU | 1 | 2.04 | SPRING | STANDARD31516 | HKLR | P |
| 11-Mar-16 | NE LANTAU | 2 | 17.97 | SPRING | STANDARD31516 | HKLR | P |
| 11-Mar-16 | NE LANTAU | 1 | 2.40 | SPRING | STANDARD31516 | HKLR | S |
| 11-Mar-16 | NE LANTAU | 2 | 6.19 | SPRING | STANDARD31516 | HKLR | S |
| 11-Mar-16 | NE LANTAU | 3 | 2.20 | SPRING | STANDARD31516 | HKLR | S |
| 22-Mar-16 | NE LANTAU | 2 | 7.42 | SPRING | STANDARD31516 | HKLR | P |
| 22-Mar-16 | NE LANTAU | 3 | 27.44 | SPRING | STANDARD31516 | HKLR | P |
| 22-Mar-16 | NE LANTAU | 4 | 2.30 | SPRING | STANDARD31516 | HKLR | P |
| 22-Mar-16 | NE LANTAU | 2 | 5.86 | SPRING | STANDARD31516 | HKLR | S |
| 22-Mar-16 | NE LANTAU | 3 | 8.18 | SPRING | STANDARD31516 | HKLR | S |
| 22-Mar-16 | NE LANTAU | 4 | 0.40 | SPRING | STANDARD31516 | HKLR | S |
| 22-Mar-16 | NW LANTAU | 2 | 3.59 | SPRING | STANDARD31516 | HKLR | P |
| 22-Mar-16 | NW LANTAU | 3 | 9.39 | SPRING | STANDARD31516 | HKLR | P |
| 22-Mar-16 | NW LANTAU | 4 | 8.10 | SPRING | STANDARD31516 | HKLR | P |
| 22-Mar-16 | NW LANTAU | 5 | 2.40 | SPRING | STANDARD31516 | HKLR | P |
| 22-Mar-16 | NW LANTAU | 2 | 1.40 | SPRING | STANDARD31516 | HKLR | S |
| 22-Mar-16 | NW LANTAU | 3 | 5.12 | SPRING | STANDARD31516 | HKLR | S |
| 23-Mar-16 | NW LANTAU | 2 | 27.12 | SPRING | STANDARD31516 | HKLR | P |
| 23-Mar-16 | NW LANTAU | 3 | 22.69 | SPRING | STANDARD31516 | HKLR | P |
| 23-Mar-16 | NW LANTAU | 2 | 4.11 | SPRING | STANDARD31516 | HKLR | S |
| 23-Mar-16 | NW LANTAU | 3 | 5.20 | SPRING | STANDARD31516 | HKLR | S |
| 5-Apr-16 | NW LANTAU | 0 | 0.83 | SPRING | STANDARD31516 | HKLR | P |
| 5-Apr-16 | NW LANTAU | 1 | 5.38 | SPRING | STANDARD31516 | HKLR | P |
| 5-Apr-16 | NW LANTAU | 2 | 21.07 | SPRING | STANDARD31516 | HKLR | P |
| 5-Apr-16 | NW LANTAU | 3 | 13.64 | SPRING | STANDARD31516 | HKLR | P |
| 5-Apr-16 | NW LANTAU | 2 | 3.00 | SPRING | STANDARD31516 | HKLR | S |
| 5-Apr-16 | NW LANTAU | 3 | 10.08 | SPRING | STANDARD31516 | HKLR | S |
| 5-Apr-16 | NE LANTAU | 1 | 1.60 | SPRING | STANDARD31516 | HKLR | P |
| 5-Apr-16 | NE LANTAU | 2 | 15.44 | SPRING | STANDARD31516 | HKLR | P |
| 5-Apr-16 | NE LANTAU | 1 | 2.10 | SPRING | STANDARD31516 | HKLR | S |
| 5-Apr-16 | NE LANTAU | 2 | 8.06 | SPRING | STANDARD31516 | HKLR | S |
| 12-Apr-16 | NE LANTAU | 2 | 3.81 | SPRING | STANDARD31516 | HKLR | P |
| 12-Apr-16 | NE LANTAU | 3 | 13.73 | SPRING | STANDARD31516 | HKLR | P |
| 12-Apr-16 | NE LANTAU | 4 | 2.60 | SPRING | STANDARD31516 | HKLR | P |
| 12-Apr-16 | NE LANTAU | 2 | 4.20 | SPRING | STANDARD31516 | HKLR | S |
| 12-Apr-16 | NE LANTAU | 3 | 6.46 | SPRING | STANDARD31516 | HKLR | S |
| 12-Apr-16 | NW LANTAU | 3 | 4.57 | SPRING | STANDARD31516 | HKLR | P |
| 12-Apr-16 | NW LANTAU | 4 | 25.36 | SPRING | STANDARD31516 | HKLR | P |
| 12-Apr-16 | NW LANTAU | 5 | 1.90 | SPRING | STANDARD31516 | HKLR | P |

Appendix I. (cont'd)

(Abbreviations: BEAU = Beaufort Sea State; P = Primary Line Effort; S = Secondary Line Effort)

| DATE | AREA | BEAU | EFFORT | SEASON | VESSEL | TYPE | P/S |
|-----------|-----------|------|--------|--------|---------------|------|-----|
| 12-Apr-16 | NW LANTAU | 3 | 5.97 | SPRING | STANDARD31516 | HKLR | S |
| 12-Apr-16 | NW LANTAU | 4 | 2.10 | SPRING | STANDARD31516 | HKLR | S |
| 15-Apr-16 | NW LANTAU | 2 | 5.14 | SPRING | STANDARD31516 | HKLR | P |
| 15-Apr-16 | NW LANTAU | 3 | 20.36 | SPRING | STANDARD31516 | HKLR | P |
| 15-Apr-16 | NW LANTAU | 4 | 6.20 | SPRING | STANDARD31516 | HKLR | P |
| 15-Apr-16 | NW LANTAU | 2 | 3.40 | SPRING | STANDARD31516 | HKLR | S |
| 15-Apr-16 | NW LANTAU | 3 | 3.10 | SPRING | STANDARD31516 | HKLR | S |
| 15-Apr-16 | NW LANTAU | 4 | 1.40 | SPRING | STANDARD31516 | HKLR | S |
| 15-Apr-16 | NE LANTAU | 2 | 14.06 | SPRING | STANDARD31516 | HKLR | P |
| 15-Apr-16 | NE LANTAU | 3 | 6.93 | SPRING | STANDARD31516 | HKLR | P |
| 15-Apr-16 | NE LANTAU | 2 | 7.11 | SPRING | STANDARD31516 | HKLR | S |
| 15-Apr-16 | NE LANTAU | 3 | 2.90 | SPRING | STANDARD31516 | HKLR | S |
| 19-Apr-16 | NE LANTAU | 3 | 10.81 | SPRING | STANDARD31516 | HKLR | P |
| 19-Apr-16 | NE LANTAU | 4 | 6.46 | SPRING | STANDARD31516 | HKLR | P |
| 19-Apr-16 | NE LANTAU | 3 | 10.03 | SPRING | STANDARD31516 | HKLR | S |
| 19-Apr-16 | NW LANTAU | 2 | 6.79 | SPRING | STANDARD31516 | HKLR | P |
| 19-Apr-16 | NW LANTAU | 3 | 15.26 | SPRING | STANDARD31516 | HKLR | P |
| 19-Apr-16 | NW LANTAU | 4 | 9.20 | SPRING | STANDARD31516 | HKLR | P |
| 19-Apr-16 | NW LANTAU | 5 | 9.70 | SPRING | STANDARD31516 | HKLR | P |
| 19-Apr-16 | NW LANTAU | 6 | 1.30 | SPRING | STANDARD31516 | HKLR | P |
| 19-Apr-16 | NW LANTAU | 2 | 3.83 | SPRING | STANDARD31516 | HKLR | S |
| 19-Apr-16 | NW LANTAU | 3 | 3.01 | SPRING | STANDARD31516 | HKLR | S |
| 19-Apr-16 | NW LANTAU | 4 | 6.39 | SPRING | STANDARD31516 | HKLR | S |
| 3-May-16 | NE LANTAU | 2 | 15.29 | SPRING | STANDARD31516 | HKLR | P |
| 3-May-16 | NE LANTAU | 3 | 1.40 | SPRING | STANDARD31516 | HKLR | P |
| 3-May-16 | NE LANTAU | 2 | 10.01 | SPRING | STANDARD31516 | HKLR | S |
| 3-May-16 | NW LANTAU | 2 | 16.24 | SPRING | STANDARD31516 | HKLR | P |
| 3-May-16 | NW LANTAU | 3 | 23.50 | SPRING | STANDARD31516 | HKLR | P |
| 3-May-16 | NW LANTAU | 2 | 7.16 | SPRING | STANDARD31516 | HKLR | S |
| 3-May-16 | NW LANTAU | 3 | 5.60 | SPRING | STANDARD31516 | HKLR | S |
| 12-May-16 | NW LANTAU | 2 | 15.26 | SPRING | STANDARD31516 | HKLR | P |
| 12-May-16 | NW LANTAU | 3 | 16.74 | SPRING | STANDARD31516 | HKLR | P |
| 12-May-16 | NW LANTAU | 2 | 7.60 | SPRING | STANDARD31516 | HKLR | S |
| 12-May-16 | NE LANTAU | 2 | 6.52 | SPRING | STANDARD31516 | HKLR | P |
| 12-May-16 | NE LANTAU | 3 | 13.33 | SPRING | STANDARD31516 | HKLR | P |
| 12-May-16 | NE LANTAU | 2 | 4.72 | SPRING | STANDARD31516 | HKLR | S |
| 12-May-16 | NE LANTAU | 3 | 6.69 | SPRING | STANDARD31516 | HKLR | S |
| 17-May-16 | NE LANTAU | 2 | 10.20 | SPRING | STANDARD31516 | HKLR | P |
| 17-May-16 | NE LANTAU | 3 | 9.92 | SPRING | STANDARD31516 | HKLR | P |
| 17-May-16 | NE LANTAU | 2 | 6.30 | SPRING | STANDARD31516 | HKLR | S |
| 17-May-16 | NE LANTAU | 3 | 4.38 | SPRING | STANDARD31516 | HKLR | S |
| 17-May-16 | NW LANTAU | 2 | 2.74 | SPRING | STANDARD31516 | HKLR | P |
| 17-May-16 | NW LANTAU | 3 | 28.07 | SPRING | STANDARD31516 | HKLR | P |
| 17-May-16 | NW LANTAU | 4 | 0.79 | SPRING | STANDARD31516 | HKLR | P |
| 17-May-16 | NW LANTAU | 3 | 7.80 | SPRING | STANDARD31516 | HKLR | S |
| 26-May-16 | NW LANTAU | 2 | 14.13 | SPRING | STANDARD31516 | HKLR | P |
| 26-May-16 | NW LANTAU | 3 | 26.67 | SPRING | STANDARD31516 | HKLR | P |
| 26-May-16 | NW LANTAU | 2 | 7.10 | SPRING | STANDARD31516 | HKLR | S |
| 26-May-16 | NW LANTAU | 3 | 6.00 | SPRING | STANDARD31516 | HKLR | S |
| 26-May-16 | NE LANTAU | 2 | 2.62 | SPRING | STANDARD31516 | HKLR | P |
| 26-May-16 | NE LANTAU | 3 | 14.38 | SPRING | STANDARD31516 | HKLR | P |
| 26-May-16 | NE LANTAU | 2 | 3.70 | SPRING | STANDARD31516 | HKLR | S |
| 26-May-16 | NE LANTAU | 3 | 6.10 | SPRING | STANDARD31516 | HKLR | S |

Appendix II. HKLR03 Chinese White Dolphin Sighting Database (March-May 2016)

(Abberviations: STG# = Sighting Number; HRD SZ = Dolphin Herd Size; BEAU = Beaufort Sea State; PSD = Perpendicular Distance; BOAT ASSOC. = Fishing Boat Association P/S: Sighting Made on Primary/Secondary Line)

| DATE | STG # | TIME | HRD SZ | AREA | BEAU | PSD | EFFORT | TYPE | NORTHING | EASTING | SEASON | BOAT ASSOC. | P/S |
|-----------|-------|------|--------|-----------|------|-----|--------|------|----------|---------|--------|-------------|-----|
| 11-Mar-16 | 1 | 1300 | 1 | NW LANTAU | 2 | ND | OFF | HKLR | 821158 | 812895 | SPRING | NONE | |
| 23-Mar-16 | 1 | 1338 | 3 | NW LANTAU | 2 | 5 | ON | HKLR | 828123 | 806459 | SPRING | NONE | P |
| 5-Apr-16 | 1 | 1059 | 8 | NW LANTAU | 2 | 454 | ON | HKLR | 824938 | 804702 | SPRING | NONE | P |
| 19-Apr-16 | 1 | 1426 | 2 | NW LANTAU | 2 | ND | OFF | HKLR | 828998 | 806471 | SPRING | NONE | |
| 19-Apr-16 | 2 | 1451 | 2 | NW LANTAU | 2 | ND | OFF | HKLR | 829109 | 806461 | SPRING | NONE | |
| 19-Apr-16 | 3 | 1504 | 3 | NW LANTAU | 2 | 177 | ON | HKLR | 829696 | 806297 | SPRING | NONE | P |
| 19-Apr-16 | 4 | 1519 | 3 | NW LANTAU | 2 | 465 | ON | HKLR | 829442 | 806050 | SPRING | NONE | S |

Appendix III. Individual dolphins identified during HKLR03 monitoring surveys in March-May 2016

| ID# | DATE | STG# | AREA |
|------------|-------------|-------------|-------------|
| CH65 | 05/04/16 | 1 | NW LANTAU |
| NL48 | 11/03/16 | 1 | NW LANTAU |
| | 05/04/16 | 1 | NW LANTAU |
| NL120 | 05/04/16 | 1 | NW LANTAU |
| NL123 | 05/04/16 | 1 | NW LANTAU |
| NL145 | 05/04/16 | 1 | NW LANTAU |
| NL202 | 19/04/16 | 1 | NW LANTAU |
| NL224 | 05/04/16 | 1 | NW LANTAU |
| NL259 | 05/04/16 | 1 | NW LANTAU |
| NL261 | 05/04/16 | 1 | NW LANTAU |
| NL264 | 05/04/16 | 1 | NW LANTAU |
| NL285 | 23/03/16 | 1 | NW LANTAU |
| | 05/04/16 | 1 | NW LANTAU |
| NL286 | 19/04/16 | 1 | NW LANTAU |
| NL287 | 05/04/16 | 1 | NW LANTAU |
| NL288 | 05/04/16 | 1 | NW LANTAU |
| NL308 | 19/04/16 | 3 | NW LANTAU |
| NL320 | 23/03/16 | 1 | NW LANTAU |

Appendix IV. Sixteen individual dolphins that were identified during March-May 2016 under HKLR03 impact phase monitoring surveys



Appendix IV. (cont'd)

NL145



NL202



NL224



NL259




Appendix IV. (cont'd)









Appendix IV. (cont'd)



Appendix V. Ranging patterns (95% kernel ranges) of 16 individual dolphins that were sighted during HKLR03 impact phase monitoring period (note: yellow dots indicates sightings made in March-May 2016)



Appendix V. (cont'd)



Appendix V. (cont'd)

