

CONTRACT NO. HY/2012/08

**Hong Kong-Zhuhai-Macao Bridge Tuen Mun – Chek Lap Kok Link
(Northern Connection Sub-sea Tunnel Section)
Dolphin Quarterly Monitoring**

*3rd Quarterly Progress Report (June-August 2014)
submitted to Dragages – Bouygues Joint Venture & ERM Hong Kong Ltd.*

Submitted by
Samuel K.Y. Hung, Ph.D., Hong Kong Cetacean Research Project

27 October 2014

1. Introduction

- 1.1. As part of the Hong Kong-Zhuhai-Macao Bridge, the Tuen Mun-Chek Lap Kok Link (TM-CLKL) Northern Connection Sub-sea Tunnel Section (Contract no. HY/2012/08) comprises the sub-sea TBM tunnels (two tubes with cross passages) across the Urmston Road to connect Tuen Area 40 and Hong Kong Boundary Crossing Facilities (HKBCF) of approximately 4 km in length with dual 2-lane carriageway, the tunnels at both the southern landfall and the northern landfall for construction of approach roads to the sub-sea TBM tunnels of approximately 1.5 km in length, as well as the northern landfall reclamation of approximately 16.5 hectares and about 20.km long seawalls. Dragages – Bouygues Joint Venture (hereinafter called the “Contractor”) was awarded as the main contractor for the Northern Connection Sub-sea Tunnel Section, and ERM Hong Kong Limited would serve as the Environmental Team to implement the Environmental Monitoring and Audit (EM&A) programme.
- 1.2. According to the updated EM&A Manual (for TM-CLKL), monthly line-transect vessel surveys for Chinese White Dolphin should be conducted to cover the Northwest (NWL) and Northeast Lantau (NEL) survey areas as in AFCD annual marine mammal monitoring programme. However, as such surveys have been undertaken by the HKLR03 and HKBCF projects in the same areas (i.e. NWL and NEL), a combined monitoring approach is recommended by the Highways Department, that the TM-CLKL EM&A project can utilize the monitoring data collected by HKLR03 or HKBCF project to avoid any redundancy in monitoring effort. Such exemption for the dolphin monitoring will end upon the completion of the dolphin monitoring carried out by HKLR03 contract.
- 1.3. In November 2013, the Director of Hong Kong Cetacean Research Project (HKCRP), Dr. Samuel Hung, has been appointed by ERM Hong Kong Limited as the dolphin specialist for the TM-CLKL Northern Connection Sub-sea Tunnel Section EM&A project. He is responsible for the dolphin monitoring study, including the data collection on Chinese White

HK CETACEAN RESEARCH PROJECT

香港鯨豚研究計劃

Dolphins during the construction phase (i.e. impact period) of the TM-CLKL project in Northwest Lantau (NWL) and Northeast Lantau (NEL) survey areas.

- 1.4. During the construction period of HKLR, the dolphin specialist would be in charge of reviewing and collating information collected by HKLR03 dolphin monitoring programme to examine any potential impacts of TM-CLKL construction works on the dolphins.
- 1.5. From the monitoring results, any changes in dolphin occurrence within the study area will be examined for possible causes, and appropriate actions and additional mitigation measures will be recommended as necessary.
- 1.6. This report is the second quarterly progress report under the TM-CLKL construction phase dolphin monitoring programme submitted to the Contractor, summarizing the results of the surveys findings during the period of June to August 2014 utilizing the survey data collected by HKLR03 project.

2. Monitoring Methodology

2.1. Vessel-based Line-transect Survey

- 2.1.1. According to the requirement of the updated EM&A manual, dolphin monitoring programme should cover all transect lines in NEL and NWL survey areas (see Figure 1) twice per month throughout the entire construction period. The co-ordinates of all transect lines conducted during the HKLR03 dolphin monitoring surveys are shown in Table 1.

Table 1 Co-ordinates of transect lines conducted by HKLR03 project

Line No.		Easting	Northing		Line No.		Easting	Northing
1	Start Point	804671	814577		13	Start Point	816506	819480
1	End Point	804671	831404		13	End Point	816506	824859
2	Start Point	805475	815457		14	Start Point	817537	820220
2	End Point	805477	826654		14	End Point	817537	824613
3	Start Point	806464	819435		15	Start Point	818568	820735
3	End Point	806464	822911		15	End Point	818568	824433
4	Start Point	807518	819771		16	Start Point	819532	821420
4	End Point	807518	829230		16	End Point	819532	824209
5	Start Point	808504	820220		17	Start Point	820451	822125
5	End Point	808504	828602		17	End Point	820451	823671
6	Start Point	809490	820466		18	Start Point	821504	822371
6	End Point	809490	825352		18	End Point	821504	823761
7	Start Point	810499	820690		19	Start Point	822513	823268
7	End Point	810499	824613		19	End Point	822513	824321

HK CETACEAN RESEARCH PROJECT 香港鯨豚研究計劃

8	Start Point	811508	820847		20	Start Point	823477	823402
8	End Point	811508	824254		20	End Point	823477	824613
9	Start Point	812516	820892		21	Start Point	805476	827081
9	End Point	812516	824254		21	End Point	805476	830562
10	Start Point	813525	820872		22	Start Point	806464	824033
10	End Point	813525	824657		22	End Point	806464	829598
11	Start Point	814556	818449		23	Start Point	814559	821739
11	End Point	814556	820992		23	End Point	814559	824768
12	Start Point	815542	818807					
12	End Point	815542	824882					

- 2.1.2. The HKLR03 survey team used standard line-transect methods (Buckland et al. 2001) to conduct the systematic vessel surveys, and followed the same technique of data collection that has been adopted over the last 16 years of marine mammal monitoring surveys in Hong Kong developed by HKCRP (see Hung 2013, 2014). For each monitoring vessel survey, a 15-m inboard vessel with an open upper deck (about 4.5 m above water surface) was used to make observations from the flying bridge area.
- 2.1.3. Two experienced observers (a data recorder and a primary observer) made up the on-effort survey team, and the survey vessel transited different transect lines at a constant speed of 13-15 km per hour. The data recorder searched with unaided eyes and filled out the datasheets, while the primary observer searched for dolphins and porpoises continuously through 7 x 50 *Fujinon* marine binoculars. Both observers searched the sea ahead of the vessel, between 270° and 90° (in relation to the bow, which is defined as 0°). One to two additional experienced observers were available on the boat to work in shift (i.e. rotate every 30 minutes) in order to minimize fatigue of the survey team members. All observers were experienced in small cetacean survey techniques and identifying local cetacean species.
- 2.1.4. During on-effort survey periods, the survey team recorded effort data including time, positions (latitude and longitude), weather conditions (Beaufort sea state and visibility), and distance traveled in each series (a continuous period of search effort) with the assistance of a handheld GPS (*Garmin eTrex Legend*).
- 2.1.5. Data including time, position and vessel speed were also automatically and continuously logged by handheld GPS throughout the entire survey for subsequent review.
- 2.1.6. When dolphins were sighted, the survey team would end the survey effort, and immediately record the initial sighting distance and angle of the dolphin group from the survey vessel, as well as the sighting time and position. Then the research vessel was diverted from its course to approach the animals for species identification, group size estimation, assessment of group composition, and behavioural observations. The perpendicular distance (PSD) of the dolphin group to the transect line was later calculated from the initial sighting distance and angle.

- 2.1.7. Survey effort being conducted along the parallel transect lines that were perpendicular to the coastlines (as indicated in Figure 1) was labeled as “primary” survey effort, while the survey effort conducted along the connecting lines between parallel lines was labeled as “secondary” survey effort. According to HKCRP long-term dolphin monitoring data, encounter rates of Chinese white dolphins deduced from effort and sighting data collected along primary and secondary lines were similar in NEL and NWL survey areas. Therefore, both primary and secondary survey effort were presented as on-effort survey effort in this report.

2.2. *Photo-identification Work*

- 2.2.1. When a group of Chinese White Dolphins were sighted during the line-transect survey, the HKLR03 survey team would end effort and approach the group slowly from the side and behind to take photographs of them. Every attempt was made to photograph every dolphin in the group, and even photograph both sides of the dolphins, since the colouration and markings on both sides may not be symmetrical.
- 2.2.2. A professional digital camera (*Canon EOS 7D* or *60D* model), equipped with long telephoto lenses (100-400 mm zoom), were available on board for researchers to take sharp, close-up photographs of dolphins as they surfaced. The images were shot at the highest available resolution and stored on Compact Flash memory cards for downloading onto a computer.
- 2.2.3. All digital images taken in the field were first examined, and those containing potentially identifiable individuals were sorted out. These photographs would then be examined in greater detail, and were carefully compared to the existing Chinese White Dolphin photo-identification catalogue maintained by HKCRP since 1995.
- 2.2.4. Chinese White Dolphins can be identified by their natural markings, such as nicks, cuts, scars and deformities on their dorsal fin and body, and their unique spotting patterns were also used as secondary identifying features (Jefferson 2000).
- 2.2.5. All photographs of each individual were then compiled and arranged in chronological order, with data including the date and location first identified (initial sighting), re-sightings, associated dolphins, distinctive features, and age classes entered into a computer database.

2.3. *Data Analysis*

- 2.3.1. Distribution Analysis – The line-transect survey data was integrated with the Geographic Information System (GIS) in order to visualize and interpret different spatial and temporal patterns of dolphin distribution using sighting positions. Location data of dolphin groups were plotted on map layers of Hong Kong using a desktop GIS (ArcView[®] 3.1) to examine their distribution patterns in details. The dataset was also stratified into different subsets to examine distribution patterns of dolphin groups with different categories of group sizes, young calves and activities.

- 2.3.2. Encounter rate analysis – Encounter rates of Chinese white dolphins (number of on-effort sightings per 100 km of survey effort, and total number of dolphins sighted on-effort per 100 km of survey effort) were calculated in NEL and NWL survey areas in relation to the amount of survey effort conducted during each month of monitoring survey. Only data collect under Beaufort 3 or below condition would be used for the encounter rate analyses. Dolphin encounter rates were calculated in two ways for comparisons with the HZMB baseline monitoring results as well as to AFCD long-term marine mammal monitoring results.

Firstly, for the comparison with the HZMB baseline monitoring results, the encounter rates were calculated using primary survey effort alone. The average encounter rate of sightings (STG) and average encounter rate of dolphins (ANI) were deduced based on the encounter rates from six events during the present quarter (i.e. six sets of line-transect surveys in North Lantau), which was also compared with the one deduced from the six events during the baseline period (i.e. six sets of line-transect surveys in North Lantau).

Secondly, the encounter rates were calculated using both primary and secondary survey effort collected under Beaufort 3 or below condition as in AFCD long-term monitoring study. The encounter rate of sightings and dolphins were deduced by dividing the total number of on-effort sightings (STG) and total number of dolphins (ANI) by the amount of survey effort for the quarterly period of June to August 2014.

- 2.3.3. Quantitative grid analysis on habitat use – To conduct quantitative grid analysis of habitat use, positions of on-effort sightings of Chinese White Dolphins collected during the quarterly impact phase monitoring period were plotted onto 1-km² grids among NWL and NEL survey areas on GIS. Sighting densities (number of on-effort sightings per km²) and dolphin densities (total number of dolphins from on-effort sightings per km²) were then calculated for each 1 km by 1 km grid with the aid of GIS. Sighting density grids and dolphin density grids were then further normalized with the amount of survey effort conducted within each grid. The total amount of survey effort spent on each grid was calculated by examining the survey coverage on each line-transect survey to determine how many times the grid was surveyed during the study period. For example, when the survey boat traversed through a specific grid 50 times, 50 units of survey effort were counted for that grid. With the amount of survey effort calculated for each grid, the sighting density and dolphin density of each grid were then normalized (i.e. divided by the unit of survey effort).

The newly-derived unit for sighting density was termed SPSE, representing the number of on-effort sightings per 100 units of survey effort. In addition, the derived unit for actual dolphin density was termed DPSE, representing the number of dolphins per 100 units of survey effort. Among the 1-km² grids that were partially covered by land, the percentage of sea area was calculated using GIS tools, and their SPSE and DPSE values were adjusted accordingly. The following formulae were used to estimate SPSE and DPSE in each 1-km² grid within the study area:

$$SPSE = ((S / E) \times 100) / SA\%$$

$$DPSE = ((D / E) \times 100) / SA\%$$

where S = total number of on-effort sightings
 D = total number of dolphins from on-effort sightings
 E = total number of units of survey effort
 SA% = percentage of sea area

- 2.3.4. Behavioural analysis – When dolphins were sighted during vessel surveys, their behaviour was observed. Different activities were categorized (i.e. feeding, socializing, traveling, and milling/resting) and recorded on sighting datasheets. This data was then input into a separate database with sighting information, which can be used to determine the distribution of behavioural data with a desktop GIS. Distribution of sightings of dolphins engaged in different activities and behaviours would then be plotted on GIS and carefully examined to identify important areas for different activities of the dolphins.
- 2.3.5. Ranging pattern analysis – Location data of individual dolphins that occurred during the 3-month impact phase monitoring period were obtained from the dolphin sighting database and photo-identification catalogue. To deduce home ranges for individual dolphins using the fixed kernel methods, the program Animal Movement Analyst Extension, was loaded as an extension with ArcView[®] 3.1 along with another extension Spatial Analyst 2.0. Using the fixed kernel method, the program calculated kernel density estimates based on all sighting positions, and provided an active interface to display kernel density plots. The kernel estimator then calculated and displayed the overall ranging area at 95% UD level.

3. Monitoring Results

3.1. *Summary of survey effort and dolphin sightings*

- 3.1.1. During the period of June to August 2014, six sets of systematic line-transect vessel surveys were conducted under the HKLR03 monitoring works to cover all transect lines in NWL and NEL survey areas twice per month.
- 3.1.2. From these HKLR03 surveys, a total of 894.40 km of survey effort was collected, with 93.6% of the total survey effort being conducted under favourable weather conditions (i.e. Beaufort Sea State 3 or below with good visibility). Among the two areas, 343.21 km and 551.19 km of survey effort were conducted in NEL and NWL survey areas respectively.
- 3.1.3. The total survey effort conducted on primary lines was 647.96 km, while the effort on secondary lines was 246.44 km. Both survey effort conducted on primary and secondary lines were considered as on-effort survey data. Summary table of the survey effort is shown in Appendix I.
- 3.1.4. During the six sets of HKLR03 monitoring surveys from June to August 2014, a total of 28 groups of 96 Chinese White Dolphins were sighted. All except two dolphin sightings were made during on-effort search. Twenty on-effort sightings were made on primary

lines, while another six on-effort sightings were made on secondary lines. In this quarterly period, almost all dolphin groups were sighted in NWL, with the exception of one group of four dolphins being sighted in NEL. Summary table of the dolphin sightings is shown in Appendix II.

3.2. *Distribution*

- 3.2.1. Distribution of dolphin sightings made during the HKLR03 monitoring surveys in June to August 2014 is shown in Figure 1. The majority of dolphin sightings were made in the western end of the North Lantau region, with higher concentration within and adjacent to the Sha Chau and Lung Kwu Chau Marine Park (Figure 1). Other dolphin sightings were scattered to the west and northeast of airport platform. The lone sighting made in NEL was located to the north of Yam O at the eastern end of the survey area (Figure 1).
- 3.2.2. Notably, none of the dolphin groups were sighted in the vicinity of TMCLKL northern landfall or southern viaduct section, and the HKLR03/HKBCF reclamation site (Figure 1).
- 3.2.3. Sighting distribution of the present impact phase monitoring period (June to August 2014) was compared to the one during the baseline monitoring period (September to November 2011). In the present quarter, dolphins have mostly avoided the NEL region, which was in stark contrast to their frequent occurrence around the Brothers Islands and in the vicinity of HKBCF reclamation site during the baseline period (Figure 1). The nearly abandonment of NEL region by the dolphins have been consistently recorded in the past six quarters.
- 3.2.4. Dolphin occurrence in the northwestern portion of North Lantau region was somewhat different between the baseline and impact phase quarters. During the present impact monitoring period, there appeared to be much fewer dolphins occurred in the middle portion of North Lantau region than during the baseline period, where dolphins supposedly moved between their core areas around Lung Kwu Chau and the Brothers Islands (Figure 1). Moreover, more dolphins were sighted between Black Point and Lung Kwu Chau during the baseline period than during the present impact monitoring period (Figure 1). A number of dolphin sightings were made to the west of Chek Lap Kok airport (especially near the HKLR09 alignment) during the baseline period, but only two sightings were made there during the present impact phase period.
- 3.2.5. As the baseline monitoring period was in the autumn season while the present monitoring period was in the summer season, a direct comparison in dolphin distribution between the two quarterly periods of summer months in 2013 and 2014 was also made to avoid the potential bias contributed by seasonal variation in distribution (Figure 2).
- 3.2.6. Among the two summer periods, only one dolphin sighting was made in NEL in the summer of 2014, while there were five sightings made there in the summer of 2013. Moreover, a lot more dolphin sightings were made in the middle and western portions of North Lantau waters (especially near Black Point, Pillar Point, to the north of airport platform and near the HKLR09 alignment) in the summer of 2013 than in the summer of 2014.

HK CETACEAN RESEARCH PROJECT

香港鯨豚研究計劃

3.3. Encounter rate

3.3.1. During the present quarterly period, the encounter rates of Chinese White Dolphins deduced from the survey effort and on-effort sighting data from the primary transect lines under favourable conditions (Beaufort 3 or below) for each set of the HKLR03 surveys in NEL and NWL are shown in Table 2. The average encounter rates deduced from the six sets of HKLR03 surveys were also compared with the ones deduced from the baseline monitoring period (September – November 2011) (Table 3).

Table 2. Dolphin encounter rates (sightings per 100 km of survey effort) during June – August 2014 deduced from HKLR03 monitoring surveys

SURVEY AREA	HKLR03 DOLPHIN MONITORING DATES	Encounter rate (STG) (no. of on-effort dolphin sightings per 100 km of survey effort)	Encounter rate (ANI) (no. of dolphins from all on-effort sightings per 100 km of survey effort)
		Primary Lines Only	Primary Lines Only
Northeast Lantau	Set 1 (3 & 5 Jun 2014)	0.00	0.00
	Set 2 (10 & 16 Jun 2014)	0.00	0.00
	Set 3 (3, 9 & 10 Jul 2014)	2.54	10.16
	Set 4 (14 & 21 Jul 2014)	0.00	0.00
	Set 5 (5 & 6 Aug 2014)	0.00	0.00
	Set 6 (15 & 19 Aug 2014)	0.00	0.00
Northwest Lantau	Set 1 (3 & 5 Jun 2014)	1.67	5.00
	Set 2 (10 & 16 Jun 2014)	0.00	0.00
	Set 3 (3, 9 & 10 Jul 2014)	3.03	10.61
	Set 4 (14 & 21 Jul 2014)	8.40	26.60
	Set 5 (5 & 6 Aug 2014)	5.63	22.52
	Set 6 (15 & 19 Aug 2014)	9.70	40.40

Table 3. Comparison of average dolphin encounter rates from impact monitoring period (June – August 2014) and baseline monitoring period (September – November 2011) (Note: encounter rates deduced from the baseline monitoring period have been recalculated based only on survey effort and on-effort sighting data made along the primary transect lines under favourable conditions)

	Encounter rate (STG) (no. of on-effort dolphin sightings per 100 km of survey effort)		Encounter rate (ANI) (no. of dolphins from all on-effort sightings per 100 km of survey effort)	
	June - August 2014	September - November 2011	June - August 2014	September - November 2011
Northeast Lantau	0.42 ± 1.04	6.00 ± 5.05	1.69 ± 4.15	22.19 ± 26.81
Northwest Lantau	4.74 ± 3.84	9.85 ± 5.85	17.52 ± 15.12	44.66 ± 29.85

3.3.2. To facilitate the comparison with the AFCD long-term monitoring results, the encounter

HK CETACEAN RESEARCH PROJECT

香港鯨豚研究計劃

rates were also calculated for the present quarter using both primary and secondary survey effort. The encounter rates of sightings (STG) and dolphins (ANI) in NWL were 5.04 sightings and 17.54 dolphins per 100 km of survey effort respectively, while the encounter rates of sightings (STG) and dolphins (ANI) in NEL were 0.29 sightings and 1.17 dolphins per 100 km of survey effort respectively.

- 3.3.3. In NEL, the average dolphin encounter rates (both STG and ANI) in the present three-month impact monitoring period was only a small fraction of the baseline value (i.e. less than 10%), and such low occurrence of dolphins in NEL have been consistently recorded in the past six quarters (Table 4).

Table 4. Comparison of average dolphin encounter rates in Northeast Lantau survey area from all quarters of impact monitoring period and baseline monitoring period (September-November 2011) (Note: encounter rates deduced from the baseline monitoring period have been recalculated based only on survey effort and on-effort sighting data made along the primary transect lines under favourable conditions)

	Encounter rate (STG) (no. of on-effort dolphin sightings per 100 km of survey effort)	Encounter rate (ANI) (no. of dolphins from all on-effort sightings per 100 km of survey effort)
September-November 2011 (Baseline)	6.00 ± 5.05	22.19 ± 26.81
December 2012-February 2013 (Impact)	3.14 ± 3.21	6.33 ± 8.64
March-May 2013 (Impact)	0.42 ± 1.03	0.42 ± 1.03
June-August 2013 (Impact)	0.88 ± 1.36	3.91 ± 8.36
September-November 2013 (Impact)	1.01 ± 1.59	3.77 ± 6.49
December 2013-February 2014 (Impact)	0.45 ± 1.10	1.34 ± 3.29
March-May 2014 (Impact)	0.00	0.00
June-August 2014 (Impact)	0.42 ± 1.04	1.69 ± 4.15

- 3.3.4. It is a serious concern that dolphin occurrence in NEL in the past six quarters (0.0-1.0 for ER(STG) and 0.0-3.9 for ER(ANI)) have been exceptionally low when compared to the baseline period (Table 4). In fact, the present quarter was the seventh consecutive quarters being accessed that have triggered the Action Level under the Event and Action Plan. As discussed recently in Hung (2014), the dramatic decline in dolphin usage of NEL waters in 2012 and 2013 (including the declines in abundance, encounter rate and habitat use in NEL, as well as shifts of individual core areas and ranges away from NEL waters) was possibly related to the HZMB construction works that were commenced in 2012.
- 3.3.5. Moreover, the average dolphin encounter rates (STG and ANI) in NWL during the present impact phase monitoring period were also much lower (reductions of 52% and 61% respectively) than the ones recorded in the 3-month baseline period, indicating a noticeable decline in dolphin usage of this survey area during the present construction period. In fact, both dolphin encounter rates in summer 2014 have dropped to the lowest since the commencement of the HKLR03 dolphin monitoring (Table 5).

HK CETACEAN RESEARCH PROJECT

香港鯨豚研究計劃

Table 5. Comparison of average dolphin encounter rates in Northwest Lantau survey area from all quarters of impact monitoring period and baseline monitoring period (September-November 2011) (Note: encounter rates deduced from the baseline monitoring period have been recalculated based only on survey effort and on-effort sighting data made along the primary transect lines under favourable conditions)

	Encounter rate (STG) (no. of on-effort dolphin sightings per 100 km of survey effort)	Encounter rate (ANI) (no. of dolphins from all on-effort sightings per 100 km of survey effort)
September-November 2011 (Baseline)	9.85 ± 5.85	44.66 ± 29.85
December 2012-February 2013 (Impact)	8.36 ± 5.03	35.90 ± 23.10
March-May 2013 (Impact)	7.75 ± 3.96	24.23 ± 18.05
June-August 2013 (Impact)	6.56 ± 3.68	27.00 ± 18.71
September-November 2013 (Impact)	8.04 ± 1.10	32.48 ± 26.51
December 2013-February 2014 (Impact)	8.21 ± 2.21	32.58 ± 11.21
March-May 2014 (Impact)	6.51 ± 3.34	19.14 ± 7.19
June-August 2014 (Impact)	4.74 ± 3.84	17.52 ± 15.12

- 3.3.6. A two-way ANOVA with repeated measures and unequal sample size was conducted to examine whether there were any significant differences in the average encounter rates between the baseline and impact monitoring periods. The two variables that were examined included the two periods (baseline and impact phases) and two locations (NEL and NWL).
- 3.3.7. For the comparison between the baseline period and the present quarter (seventh quarter of the impact phase being assessed), the p-value for the differences in average dolphin encounter rates of STG and ANI were 0.0199 and 0.0597 respectively. If the alpha value is set at 0.1, significant difference was detected between the baseline and present quarters in both dolphin encounter rates of STG and ANI.
- 3.3.8. For the comparison between the baseline period and the cumulative quarters in impact phase (i.e. first seven quarters of the impact phase being assessed), the p-value for the differences in average dolphin encounter rates of STG and ANI were 0.0037 and 0.0013 respectively. Even if the alpha value is set at 0.01, significant differences were detected in both the average dolphin encounter rates of STG and ANI (i.e. between the two periods and the locations).
- 3.4. *Group size*
- 3.4.1. Group size of Chinese White Dolphins ranged from one to eight individuals per group in North Lantau region during June – August 2014. The average dolphin group sizes from these three months were compared with the ones deduced from the baseline period in September to November 2011, as shown in Table 6.

Table 6. Comparison of average dolphin group sizes from impact monitoring period (June – August 2014) and baseline monitoring period (September – November 2011)

	Average Dolphin Group Size	
	June – August 2014	September – November 2011
Overall	3.43 ± 1.95 (n = 28)	3.72 ± 3.13 (n = 66)
Northeast Lantau	4.00 ± 0.00 (n = 1)	3.18 ± 2.16 (n = 17)
Northwest Lantau	3.41 ± 1.99 (n = 27)	3.92 ± 3.40 (n = 49)

- 3.4.2. The average dolphin group sizes in the entire North Lantau region as well as in NWL waters during June – August 2014 were lower than the ones recorded during the three-month baseline period (Table 6). In fact, 17 of the 28 groups were composed of 1-3 individuals only, while no dolphin group was composed of more than 10 individuals.
- 3.4.3. Distribution of dolphins with larger group sizes (five individuals or more per group) during the present quarter is shown in Figure 3, with comparison to the one in baseline period. During the summer of 2014, distribution of all larger dolphin groups were concentrated within and around the Sha Chau and Lung Kwu Chau Marine Park (Figure 3). This distribution pattern was quite different from the baseline period, when the larger dolphin groups were distributed more evenly in NWL waters with a few more sighted in NEL waters (Figure 3).
- 3.5. *Habitat use*
- 3.5.1. From June to August 2014, the most heavily utilized habitats by Chinese White Dolphins mainly concentrated within and around the marine park area (Figures 4a and 4b). Only one grid in NEL recorded the presence of dolphins. Moreover, all grids near TMCLKL and HKLR09 alignments as well as the HKLR03/HKBCF reclamation sites did not record any presence of dolphins during on-effort search in the present quarterly period.
- 3.5.2. However, it should be emphasized that the amount of survey effort collected in each grid during the three-month period was fairly low (6-12 units of survey effort for most grids), and therefore the habitat use pattern derived from the three-month dataset should be treated with caution. A more complete picture of dolphin habitat use pattern will be presented when more survey effort for each grid will be collected throughout the impact phase monitoring programme.
- 3.5.3. When compared with the habitat use patterns during the baseline period, dolphin usage in NEL was dramatically different from the present impact monitoring period (Figure 5). During the baseline period, nine grids between Siu Mo To and Shum Shui Kok recorded moderately high to high dolphin densities, which was in stark contrast to the very rare occurrence of dolphins during the present impact phase period (Figure 5).
- 3.5.4. The density patterns between the baseline and impact phase monitoring periods were also

different in NWL, with higher dolphin usage near Black Point, as well as between Pillar Point and airport platform during the baseline period (Figure 5).

3.6. *Mother-calf pairs*

- 3.6.1. During the three-month study period, only three unspotted juveniles (UJ) were sighted in NWL survey areas. These young calves comprised of 3.1% of all animals sighted, which was much lower than the percentage recorded during the baseline monitoring period (6.8%).
- 3.6.2. The few young calves were found near Lung Kwu Chau, Sha Chau and Shum Wat (Figure 6), which was very different from their distribution pattern during the baseline period when young calves were sighted throughout the NWL survey area as well as a few sighted in NEL waters. None of these young calves were sighted in the vicinity of the HKLR09/TMCLKL alignments and HKBCF/HKLR03 reclamation sites during the present quarter (Figure 6).

3.7. *Activities and associations with fishing boats*

- 3.7.1. A total of four dolphin sightings were associated with socializing and traveling activities during the three-month study period. Notably, no feeding activity of dolphin was observed during the present quarter, which was in contrast to the relatively high percentage of feeding activities recorded during the baseline period (11.6%). On the contrary, the percentage of socializing activities during the present impact phase monitoring period (7.1%) was slightly higher than the one recorded during the baseline period (5.4%).
- 3.7.2. Distribution of dolphins engaged in socializing and traveling activities during the present three-month period is shown in Figure 7. The two sightings associated with socializing activities occurred near Sha Chau, while the two sightings associated with traveling activities were found adjacent to Lung Kwu Chau (Figure 7). Distribution of dolphin sightings associated with these activities during the impact phase was drastically different from the distribution pattern of these activities during the baseline period (Figure 7).
- 3.7.3. During the three-month period, none of the 28 dolphin groups was found to be associated with an operating fishing vessels in North Lantau waters. The rare events of fishing boat association in the present and previous quarters were consistently found, and were likely related to the recent trawl ban being implemented in December 2012 in Hong Kong waters.

3.8. *Summary of photo-identification works*

- 3.8.1. From June to August 2014, over 2,000 digital photographs of Chinese White Dolphins were taken during the impact phase monitoring surveys for the photo-identification work.
- 3.8.2. In total, 32 individuals sighted 44 times altogether were identified (see summary table in Appendix III and photographs of identified individuals in Appendix IV). All except four of these re-sightings were made in NWL. Four individuals (NL123, NL139, NL261 and NL285) were sighted once during the lone sighting made in NEL in the present quarter.

- 3.8.3. Almost all identified individuals were sighted only once or twice during the three-month period, with the exception of one individual (NL272) being sighted thrice.
- 3.8.4. Notably, 11 of these 32 individuals were also sighted in West Lantau waters during the HKLR09 monitoring surveys during the same three-month period, showing their extensive movement between North and West Lantau regions. In particular, two individuals (NL139 and NL261) were sighted in both NEL and WL during the same quarter.
- 3.8.5. Six well-recognized females (NL93, NL104, NL123, NL145, NL202 and WL124) were accompanied with their calves during their re-sightings. Most of these mothers were frequently sighted with their calves throughout the HKLR03 impact phase monitoring period since October 2012.
- 3.9. *Individual range use*
- 3.9.1. Ranging patterns of the 32 individuals identified during the three-month study period were determined by fixed kernel method, and are shown in Appendix V.
- 3.9.2. With the exception of a few individuals, most identified dolphins sighted in this quarter were utilizing their range use in NWL (and some also in WL), but have avoided the NEL waters where many of them have utilized as their core areas in the past (Appendix V). This is in contrary to the extensive movements between NEL and NWL survey areas observed in the earlier impact monitoring quarters as well as during the baseline period.
- 3.9.3. For many individuals that have previously utilized the Brothers Islands as their major core area of activities, they have apparently shifted their range use away from this important habitat (e.g. NL93, NL123, NL136, NL261; Appendix V). Such shifts of range use and core area use were also documented by Hung (2014), as well as in the past monitoring quarters in 2013 and 2014 under the present study.

4. Conclusion

- 4.1. During this quarter of dolphin monitoring, no adverse impact from the activities of the TMCLKL construction project on Chinese White Dolphins was noticeable from general observations.
- 4.2. Although the dolphins infrequently occurred along the alignment of TMCLKL northern connection sub-sea tunnel section in the past and during the baseline monitoring period, it is apparent that dolphin usage has been significantly reduced in NEL, and many individuals have shifted away from the important habitat around the Brothers Islands.
- 4.3. It is critical to monitor the dolphin usage in North Lantau region in the upcoming quarters, to determine whether the dolphins are continuously affected by the various construction activities in relation to the HZMB-related works, and whether suitable mitigation measure can be applied to revert the situation.

5. References

- Buckland, S. T., Anderson, D. R., Burnham, K. P., Laake, J. L., Borchers, D. L., and Thomas, L. 2001. Introduction to distance sampling: estimating abundance of biological populations. Oxford University Press, London.
- Hung, S. K. 2013. Monitoring of Marine Mammals in Hong Kong waters: final report (2012-13). An unpublished report submitted to the Agriculture, Fisheries and Conservation Department, 168 pp.
- Hung, S. K. 2014. Monitoring of marine mammals in Hong Kong waters – data collection: final report (2013-14). An unpublished report submitted to the Agriculture, Fisheries and Conservation Department of Hong Kong SAR Government, 231 pp.
- Jefferson, T. A. 2000. Population biology of the Indo-Pacific hump-backed dolphin in Hong Kong waters. Wildlife Monographs 144:1-65.





Figure 1. Distribution of Chinese white dolphin sighting in Northwest and Northeast Lantau during HKLR03 impact phase (top) and baseline monitoring surveys (bottom)





Figure 2. Distribution of Chinese white dolphin sighting in Northwest and Northeast Lantau during the same summer quarters of HKLR03 impact phase in 2014 (top) and 2013 (bottom)





Figure 3. Distribution of Chinese white dolphins with larger group sizes during HKLR03 impact phase (top) and baseline monitoring surveys (bottom) (green dots: group sizes of 5 or more; purple dots: group sizes of 10 or more)




Figure 4a. Sighting density of Chinese white dolphins with corrected survey effort per km² in Northeast and Northwest Lantau survey areas, using data collected during HKLR03 impact monitoring period (Jun-Aug 14) (SPSE = no. of on-effort sightings per 100 units of survey effort)




Figure 4b. Density of Chinese white dolphins with corrected survey effort per km² in Northeast and Northwest Lantau survey areas, using data collected during HKLR03 impact monitoring period (Jun-Aug 14) (DPSE = no. of dolphins per 100 units of survey effort)




Figure 5. Comparison of density of Chinese white dolphins with corrected survey effort per km² in Northwest and Northeast Lantau survey area between the impact monitoring period (June-August 2014) and baseline monitoring period (September-November 2011) (DPSE = no. of dolphins per 100 units of survey effort)





Figure 6. Distribution of young calves of Chinese white dolphins during HKLR03 impact phase (top) and baseline monitoring surveys (bottom)





Figure 7. Distribution of Chinese white dolphins engaged in feeding (purple dots), socializing (pink dots) and traveling (green dots) activities during HKLR03 impact phase (top) and baseline monitoring surveys (bottom)

Appendix I. HKLR03 Survey Effort Database (June-August 2014)

(Abbreviations: BEAU = Beaufort Sea State; P = Primary Line Effort; S = Secondary Line Effort)

DATE	AREA	BEAU	EFFORT	SEASON	VESSEL	TYPE	P/S
3-Jun-14	NE LANTAU	2	14.31	SUMMER	STANDARD31516	HKLR	P
3-Jun-14	NE LANTAU	3	2.60	SUMMER	STANDARD31516	HKLR	P
3-Jun-14	NE LANTAU	2	10.89	SUMMER	STANDARD31516	HKLR	S
3-Jun-14	NW LANTAU	2	6.52	SUMMER	STANDARD31516	HKLR	P
3-Jun-14	NW LANTAU	3	23.00	SUMMER	STANDARD31516	HKLR	P
3-Jun-14	NW LANTAU	4	10.70	SUMMER	STANDARD31516	HKLR	P
3-Jun-14	NW LANTAU	2	3.78	SUMMER	STANDARD31516	HKLR	S
3-Jun-14	NW LANTAU	3	9.70	SUMMER	STANDARD31516	HKLR	S
5-Jun-14	NE LANTAU	1	5.65	SUMMER	STANDARD31516	HKLR	P
5-Jun-14	NE LANTAU	2	10.52	SUMMER	STANDARD31516	HKLR	P
5-Jun-14	NE LANTAU	3	4.20	SUMMER	STANDARD31516	HKLR	P
5-Jun-14	NE LANTAU	1	2.20	SUMMER	STANDARD31516	HKLR	S
5-Jun-14	NE LANTAU	2	6.23	SUMMER	STANDARD31516	HKLR	S
5-Jun-14	NE LANTAU	3	2.10	SUMMER	STANDARD31516	HKLR	S
5-Jun-14	NW LANTAU	2	13.90	SUMMER	STANDARD31516	HKLR	P
5-Jun-14	NW LANTAU	3	16.56	SUMMER	STANDARD31516	HKLR	P
5-Jun-14	NW LANTAU	2	3.70	SUMMER	STANDARD31516	HKLR	S
5-Jun-14	NW LANTAU	3	3.61	SUMMER	STANDARD31516	HKLR	S
10-Jun-14	NW LANTAU	2	6.21	SUMMER	STANDARD31516	HKLR	P
10-Jun-14	NW LANTAU	3	31.70	SUMMER	STANDARD31516	HKLR	P
10-Jun-14	NW LANTAU	4	2.50	SUMMER	STANDARD31516	HKLR	P
10-Jun-14	NW LANTAU	2	9.29	SUMMER	STANDARD31516	HKLR	S
10-Jun-14	NW LANTAU	3	4.10	SUMMER	STANDARD31516	HKLR	S
10-Jun-14	NE LANTAU	2	12.34	SUMMER	STANDARD31516	HKLR	P
10-Jun-14	NE LANTAU	3	3.50	SUMMER	STANDARD31516	HKLR	P
10-Jun-14	NE LANTAU	2	10.53	SUMMER	STANDARD31516	HKLR	S
10-Jun-14	NE LANTAU	3	0.73	SUMMER	STANDARD31516	HKLR	S
16-Jun-14	NW LANTAU	2	3.11	SUMMER	STANDARD31516	HKLR	P
16-Jun-14	NW LANTAU	3	13.98	SUMMER	STANDARD31516	HKLR	P
16-Jun-14	NW LANTAU	4	14.31	SUMMER	STANDARD31516	HKLR	P
16-Jun-14	NW LANTAU	3	4.28	SUMMER	STANDARD31516	HKLR	S
16-Jun-14	NW LANTAU	4	3.43	SUMMER	STANDARD31516	HKLR	S
16-Jun-14	NE LANTAU	1	1.40	SUMMER	STANDARD31516	HKLR	P
16-Jun-14	NE LANTAU	2	18.35	SUMMER	STANDARD31516	HKLR	P
16-Jun-14	NE LANTAU	1	0.30	SUMMER	STANDARD31516	HKLR	S
16-Jun-14	NE LANTAU	2	10.55	SUMMER	STANDARD31516	HKLR	S
3-Jul-14	NE LANTAU	2	1.89	SUMMER	STANDARD31516	HKLR	P
3-Jul-14	NE LANTAU	2	2.14	SUMMER	STANDARD31516	HKLR	S
3-Jul-14	NW LANTAU	2	7.87	SUMMER	STANDARD31516	HKLR	P
3-Jul-14	NW LANTAU	3	23.09	SUMMER	STANDARD31516	HKLR	P
3-Jul-14	NW LANTAU	4	5.90	SUMMER	STANDARD31516	HKLR	P
3-Jul-14	NW LANTAU	2	2.90	SUMMER	STANDARD31516	HKLR	S
3-Jul-14	NW LANTAU	3	7.84	SUMMER	STANDARD31516	HKLR	S
3-Jul-14	NW LANTAU	4	0.60	SUMMER	STANDARD31516	HKLR	S
9-Jul-14	NW LANTAU	1	1.80	SUMMER	STANDARD31516	HKLR	P
9-Jul-14	NW LANTAU	2	9.28	SUMMER	STANDARD31516	HKLR	P
9-Jul-14	NW LANTAU	2	3.22	SUMMER	STANDARD31516	HKLR	S
10-Jul-14	NW LANTAU	1	8.81	SUMMER	STANDARD31516	HKLR	P
10-Jul-14	NW LANTAU	2	12.85	SUMMER	STANDARD31516	HKLR	P
10-Jul-14	NW LANTAU	3	2.29	SUMMER	STANDARD31516	HKLR	P
10-Jul-14	NW LANTAU	1	0.73	SUMMER	STANDARD31516	HKLR	S

Appendix I. (cont'd)

(Abbreviations: BEAU = Beaufort Sea State; P = Primary Line Effort; S = Secondary Line Effort)

DATE	AREA	BEAU	EFFORT	SEASON	VESSEL	TYPE	P/S
10-Jul-14	NW LANTAU	2	6.69	SUMMER	STANDARD31516	HKLR	S
10-Jul-14	NE LANTAU	1	14.94	SUMMER	STANDARD31516	HKLR	P
10-Jul-14	NE LANTAU	2	16.33	SUMMER	STANDARD31516	HKLR	P
10-Jul-14	NE LANTAU	3	6.20	SUMMER	STANDARD31516	HKLR	P
10-Jul-14	NE LANTAU	1	3.93	SUMMER	STANDARD31516	HKLR	S
10-Jul-14	NE LANTAU	2	6.90	SUMMER	STANDARD31516	HKLR	S
10-Jul-14	NE LANTAU	3	0.80	SUMMER	STANDARD31516	HKLR	S
14-Jul-14	NW LANTAU	2	19.59	SUMMER	STANDARD31516	HKLR	P
14-Jul-14	NW LANTAU	3	11.09	SUMMER	STANDARD31516	HKLR	P
14-Jul-14	NW LANTAU	2	2.05	SUMMER	STANDARD31516	HKLR	S
14-Jul-14	NW LANTAU	3	3.80	SUMMER	STANDARD31516	HKLR	S
14-Jul-14	NW LANTAU	4	0.93	SUMMER	STANDARD31516	HKLR	S
14-Jul-14	NE LANTAU	1	2.00	SUMMER	STANDARD31516	HKLR	P
14-Jul-14	NE LANTAU	2	14.57	SUMMER	STANDARD31516	HKLR	P
14-Jul-14	NE LANTAU	3	2.40	SUMMER	STANDARD31516	HKLR	P
14-Jul-14	NE LANTAU	4	1.20	SUMMER	STANDARD31516	HKLR	P
14-Jul-14	NE LANTAU	2	10.51	SUMMER	STANDARD31516	HKLR	S
14-Jul-14	NE LANTAU	3	0.30	SUMMER	STANDARD31516	HKLR	S
21-Jul-14	NW LANTAU	1	5.9	SUMMER	STANDARD31516	HKLR	P
21-Jul-14	NW LANTAU	2	31.1	SUMMER	STANDARD31516	HKLR	P
21-Jul-14	NW LANTAU	3	3.7	SUMMER	STANDARD31516	HKLR	P
21-Jul-14	NW LANTAU	2	7.9	SUMMER	STANDARD31516	HKLR	S
21-Jul-14	NW LANTAU	3	4.9	SUMMER	STANDARD31516	HKLR	S
21-Jul-14	NE LANTAU	1	2.8	SUMMER	STANDARD31516	HKLR	P
21-Jul-14	NE LANTAU	2	13.7	SUMMER	STANDARD31516	HKLR	P
21-Jul-14	NE LANTAU	2	10.7	SUMMER	STANDARD31516	HKLR	S
5-Aug-14	NE LANTAU	1	8.40	SUMMER	STANDARD31516	HKLR	P
5-Aug-14	NE LANTAU	2	5.80	SUMMER	STANDARD31516	HKLR	P
5-Aug-14	NE LANTAU	3	2.10	SUMMER	STANDARD31516	HKLR	P
5-Aug-14	NE LANTAU	1	6.20	SUMMER	STANDARD31516	HKLR	S
5-Aug-14	NE LANTAU	2	4.80	SUMMER	STANDARD31516	HKLR	S
5-Aug-14	NW LANTAU	1	8.00	SUMMER	STANDARD31516	HKLR	P
5-Aug-14	NW LANTAU	2	30.30	SUMMER	STANDARD31516	HKLR	P
5-Aug-14	NW LANTAU	3	1.70	SUMMER	STANDARD31516	HKLR	P
5-Aug-14	NW LANTAU	1	1.50	SUMMER	STANDARD31516	HKLR	S
5-Aug-14	NW LANTAU	2	9.90	SUMMER	STANDARD31516	HKLR	S
6-Aug-14	NW LANTAU	1	4.30	SUMMER	STANDARD31516	HKLR	P
6-Aug-14	NW LANTAU	2	21.55	SUMMER	STANDARD31516	HKLR	P
6-Aug-14	NW LANTAU	3	5.21	SUMMER	STANDARD31516	HKLR	P
6-Aug-14	NW LANTAU	1	2.30	SUMMER	STANDARD31516	HKLR	S
6-Aug-14	NW LANTAU	2	4.05	SUMMER	STANDARD31516	HKLR	S
6-Aug-14	NW LANTAU	3	0.30	SUMMER	STANDARD31516	HKLR	S
6-Aug-14	NE LANTAU	1	17.62	SUMMER	STANDARD31516	HKLR	P
6-Aug-14	NE LANTAU	2	2.26	SUMMER	STANDARD31516	HKLR	P
6-Aug-14	NE LANTAU	1	10.52	SUMMER	STANDARD31516	HKLR	S
15-Aug-14	NW LANTAU	2	7.71	SUMMER	STANDARD31516	HKLR	P
15-Aug-14	NW LANTAU	3	29.93	SUMMER	STANDARD31516	HKLR	P
15-Aug-14	NW LANTAU	3	9.92	SUMMER	STANDARD31516	HKLR	S
15-Aug-14	NW LANTAU	4	2.64	SUMMER	STANDARD31516	HKLR	S
15-Aug-14	NE LANTAU	2	17.22	SUMMER	STANDARD31516	HKLR	P
15-Aug-14	NE LANTAU	3	0.58	SUMMER	STANDARD31516	HKLR	P
15-Aug-14	NE LANTAU	2	8.54	SUMMER	STANDARD31516	HKLR	S
15-Aug-14	NE LANTAU	3	1.26	SUMMER	STANDARD31516	HKLR	S

Appendix I. (cont'd)

(Abbreviations: BEAU = Beaufort Sea State; P = Primary Line Effort; S = Secondary Line Effort)

DATE	AREA	BEAU	EFFORT	SEASON	VESSEL	TYPE	P/S
19-Aug-14	NE LANTAU	1	1.46	SUMMER	STANDARD31516	HKLR	P
19-Aug-14	NE LANTAU	2	11.20	SUMMER	STANDARD31516	HKLR	P
19-Aug-14	NE LANTAU	3	5.91	SUMMER	STANDARD31516	HKLR	P
19-Aug-14	NE LANTAU	4	0.80	SUMMER	STANDARD31516	HKLR	P
19-Aug-14	NE LANTAU	2	4.35	SUMMER	STANDARD31516	HKLR	S
19-Aug-14	NE LANTAU	3	6.48	SUMMER	STANDARD31516	HKLR	S
19-Aug-14	NW LANTAU	2	1.16	SUMMER	STANDARD31516	HKLR	P
19-Aug-14	NW LANTAU	3	23.08	SUMMER	STANDARD31516	HKLR	P
19-Aug-14	NW LANTAU	4	3.24	SUMMER	STANDARD31516	HKLR	P
19-Aug-14	NW LANTAU	5	3.69	SUMMER	STANDARD31516	HKLR	P
19-Aug-14	NW LANTAU	3	4.32	SUMMER	STANDARD31516	HKLR	S
19-Aug-14	NW LANTAU	4	7.12	SUMMER	STANDARD31516	HKLR	S

Appendix II. HKLR03 Chinese White Dolphin Sighting Database (June-August 2014)

(Abbreviations: STG# = Sighting Number; HRD SZ = Dolphin Herd Size; BEAU = Beaufort Sea State; PSD = Perpendicular Distance; BOAT ASSOC. = Fishing Boat Association; P/S: Sighting Made on Primary/Secondary Line)


DATE	STG #	TIME	HRD SZ	AREA	BEAU	PSD	EFFORT	TYPE	NORTHING	EASTING	SEASON	BOAT ASSOC.	P/S
5-Jun-14	1	1400	3	NW LANTAU	3	184	ON	HKLR	827350	805448	SUMMER	NONE	P
5-Jun-14	2	1413	3	NW LANTAU	3	20	ON	HKLR	826719	805344	SUMMER	NONE	S
16-Jun-14	1	1408	1	NW LANTAU	3	ND	OFF	HKLR	827538	805459	SUMMER	NONE	
3-Jul-14	1	958	4	NE LANTAU	2	317	ON	HKLR	823230	820459	SUMMER	NONE	P
3-Jul-14	2	1302	4	NW LANTAU	3	ND	OFF	HKLR	821327	811071	SUMMER	NONE	
3-Jul-14	3	1642	2	NW LANTAU	3	161	ON	HKLR	814628	804722	SUMMER	NONE	P
10-Jul-14	1	1110	5	NW LANTAU	2	588	ON	HKLR	827483	805459	SUMMER	NONE	P
10-Jul-14	2	1150	5	NW LANTAU	2	0	ON	HKLR	829928	806565	SUMMER	NONE	S
14-Jul-14	1	1022	3	NW LANTAU	2	572	ON	HKLR	816276	805395	SUMMER	NONE	P
14-Jul-14	2	1036	1	NW LANTAU	2	866	ON	HKLR	819222	805442	SUMMER	NONE	P
14-Jul-14	3	1044	5	NW LANTAU	2	118	ON	HKLR	820484	805434	SUMMER	NONE	P
14-Jul-14	4	1105	7	NW LANTAU	2	471	ON	HKLR	822311	805448	SUMMER	NONE	P
14-Jul-14	5	1144	2	NW LANTAU	2	819	ON	HKLR	827173	805448	SUMMER	NONE	P
21-Jul-14	1	1113	1	NW LANTAU	2	694	ON	HKLR	823509	804668	SUMMER	NONE	P
21-Jul-14	2	1436	2	NW LANTAU	2	325	ON	HKLR	821325	812267	SUMMER	NONE	S
5-Aug-14	1	1413	8	NW LANTAU	2	428	ON	HKLR	826185	806764	SUMMER	NONE	P
5-Aug-14	2	1435	4	NW LANTAU	2	0	ON	HKLR	827426	806458	SUMMER	NONE	P
5-Aug-14	3	1444	2	NW LANTAU	2	990	ON	HKLR	828943	806461	SUMMER	NONE	P
5-Aug-14	4	1515	2	NW LANTAU	2	452	ON	HKLR	827872	804667	SUMMER	NONE	P
6-Aug-14	1	1110	3	NW LANTAU	3	10	ON	HKLR	826730	805323	SUMMER	NONE	S
6-Aug-14	2	1151	1	NW LANTAU	2	17	ON	HKLR	829773	806359	SUMMER	NONE	S
15-Aug-14	1	1029	5	NW LANTAU	3	393	ON	HKLR	818936	804648	SUMMER	NONE	P
15-Aug-14	2	1041	7	NW LANTAU	3	15	ON	HKLR	821006	804652	SUMMER	NONE	P
15-Aug-14	3	1218	3	NW LANTAU	3	0	ON	HKLR	823429	806027	SUMMER	NONE	S
15-Aug-14	4	1305	2	NW LANTAU	2	749	ON	HKLR	823524	808510	SUMMER	NONE	P
15-Aug-14	5	1310	6	NW LANTAU	3	83	ON	HKLR	824321	808501	SUMMER	NONE	P
19-Aug-14	1	1338	2	NW LANTAU	3	105	ON	HKLR	825220	807514	SUMMER	NONE	P
19-Aug-14	2	1536	3	NW LANTAU	2	113	ON	HKLR	823076	805450	SUMMER	NONE	P

Appendix III. Individual dolphins identified during HKLR03 monitoring surveys in June-August 2014


ID#	DATE	STG#	AREA
CH34	10/07/14	1	NW LANTAU
NL46	05/08/14	2	NW LANTAU
NL48	05/08/14	1	NW LANTAU
	19/08/14	1	NW LANTAU
NL80	14/07/14	4	NW LANTAU
NL93	10/07/14	1	NW LANTAU
	05/08/14	1	NW LANTAU
NL104	05/08/14	1	NW LANTAU
NL123	03/07/14	1	NE LANTAU
	15/08/14	5	NW LANTAU
NL136	05/06/14	2	NW LANTAU
NL139	03/07/14	1	NE LANTAU
NL145	14/07/14	3	NW LANTAU
NL182	10/07/14	2	NW LANTAU
NL202	19/08/14	1	NW LANTAU
	19/08/14	2	NW LANTAU
NL210	10/07/14	2	NW LANTAU
NL242	05/08/14	1	NW LANTAU
NL247	14/07/14	4	NW LANTAU
	15/08/14	2	NW LANTAU
NL259	05/08/14	1	NW LANTAU
NL261	03/07/14	1	NE LANTAU
NL272	05/06/14	1	NW LANTAU
	05/06/14	2	NW LANTAU
	15/08/14	5	NW LANTAU
NL278	15/08/14	2	NW LANTAU
NL284	15/08/14	5	NW LANTAU
NL285	03/07/14	1	NE LANTAU
	15/08/14	5	NW LANTAU
NL286	15/08/14	5	NW LANTAU
	19/08/14	2	NW LANTAU

ID#	DATE	STG#	AREA
NL287	14/07/14	3	NW LANTAU
	15/08/14	5	NW LANTAU
NL300	14/07/14	4	NW LANTAU
NL301	14/07/14	4	NW LANTAU
NL307	15/08/14	5	NW LANTAU
WL28	15/08/14	2	NW LANTAU
WL30	10/07/14	1	NW LANTAU
WL46	15/08/14	2	NW LANTAU
WL124	03/07/14	3	NW LANTAU
	15/08/14	2	NW LANTAU
WL188	06/08/14	1	NW LANTAU
	15/08/14	2	NW LANTAU
WL214	15/08/14	2	NW LANTAU

Appendix IV. Thirty-two individual dolphins that were identified during June – August 2014 under HKLR03 impact phase monitoring surveys



Appendix IV. (cont'd)



Appendix IV. (cont'd)

NL139



NL145




NL182



NL202



Appendix IV. (cont'd)



Appendix IV. (cont'd)

NL261



NL272



NL278



NL284



Appendix IV. (cont'd)

NL285



NL286




NL287



NL300




Appendix IV. (cont'd)




Appendix IV. (cont'd)









Appendix V. Ranging patterns (95% kernel ranges) of 32 individual dolphins that were sighted during HKLR03 impact phase monitoring period (note: yellow dots indicates sightings made in June – August 2014)




Appendix V. (cont'd)




Appendix V. (cont'd)



Appendix V. (cont'd)



Appendix V. (cont'd)



Appendix V. (cont'd)

